For Research Use Only

TAF11 Polyclonal antibody

Catalog Number: 16114-1-AP



Purification Method:

IHC 1:50-1:500

IF 1:20-1:200

Basic Information

Catalog Number: GenBank Accession Number:

16114-1-AP BC021972 Antigen affinity purification Size: GeneID (NCBI): **Recommended Dilutions:**

WB 1:200-1:1000 150ul, Concentration: 550 µg/ml by 6882

Nanodrop and 347 µg/ml by Bradford Full Name: IP 0.5-4.0 ug for IP and 1:200-1:1000 method using BSA as the standard; TAF11 RNA polymerase II, TATA box

binding protein (TBP)-associated Rabbit factor, 28kDa Calculated MW: Isotype:

IgG 211 aa, 23 kDa Observed MW: Immunogen Catalog Number: 28 kDa AG9039

Applications

Tested Applications:

IF, IHC, IP, WB, ELISA WB: HepG2 cells, HeLa cells

Species Specificity: IP: HeLa cells.

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed

with citrate buffer pH 6.0

Positive Controls:

IHC: human colon cancer tissue,

IF: HepG2 cells,

Background Information

TAF11, also named as Transcription initiation factor TFIID 28 kDa subunit, is a 211 amino acid protein, which Belongs to the TAF11 family. TAF11 as a core TAFII present in both of the previously described TFIID species which either lack or contain TAFII30. The calcualted molecular weight of TAF11 is 23 kDa, but modified TAF11 is about 28kDa. (PMID: 10336414)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1%BSA


Selected Validation Data




IP result of anti-TAF11 (IP:16114-1-AP, 4ug; Detection:16114-1-AP 1:300) with HeLa cells lysate 1600 ug.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 16114-1-AP (TAF11 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HepG2 cells were subjected to SDS PAGE followed by western blot with 16114-1-AP (TAF11 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells using 16114-1-AP (TAF11 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).