For Research Use Only

TAF11 Polyclonal antibody

Catalog Number:16114-1-AP




Basic Information	Catalog Number: 16114-1-AP	GenBank Accession Number: BC021972		Purification Method: Antigen affinity purification	
	Size: 150ul, Concentration: 550 ug/ml by Nanodrop and 347 ug/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG9039	GeneID (NCBI): 6882 UNIPROT ID: Q15544 Full Name: TAF 11 RNA polymerase binding protein (TBP)-ar factor, 28kDa Calculated MW: 211 aa, 23 kDa Observed MW: 28 kDa		Recommended Dilutions: WB 1:200-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500 IF/ICC 1:20-1:200	
Applications	Tested Applications: WB, IHC, IF/ICC, IP, ELISA				
	Species Specificity: human, mouse, rat		WB : HepG2 cells, HeLa cells IP : HeLa cells,		
	Note-IHC: suggested antigen re	etrieval with	IHC : human co	nan colon cancer tissue,	
	TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		IF/ICC : HepG	2 cells,	
Background Information	TAF11, also named as Transcription initiation factor TFIID 28 kDa subunit, is a 211 amino acid protein, which Belongs to the TAF11 family. TAF11 as a core TAFII present in both of the previously described TFIID species which either lack or contain TAFII30. The calcualted molecular weight of TAF11 is 23 kDa, but modified TAF11 is about 28kDa. (PMID: 10336414)				
Storage	Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°				
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	0, 1			


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




IP result of anti-TAF11 (IP:16114-1-AP, 4ug; Detection:16114-1-AP 1:300) with HeLa cells lysate 1600 ug.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 16114-1-AP (TAF11 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HepG2 cells were subjected to SDS PAGE followed by western blot with 16114-1-AP (TAF11 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of HepG2 cells using 16114-1-AP (TAF11 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).