### For Research Use Only

# RAC1/2/3 Polyclonal antibody

Catalog Number:16117-1-AP

**Featured Product** 

8 Publications



**Basic Information** 

Catalog Number:

16117-1-AP

GenBank Accession Number:

BC015197

21 kDa

GeneID (NCBI): Size: 150ul, Concentration: 500 µg/ml by 5881

Nanodrop and 300 µg/ml by Bradford Full Name:

method using BSA as the standard;

ras-related C3 botulinum toxin

substrate 3 (rho family, small GTP Rabbit binding protein Rac3)

Calculated MW: Isotype: 192 aa, 21 kDa IgG Immunogen Catalog Number: Observed MW:

AG9045

IP, WB, ELISA

human, mouse, rat

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:10000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB

**Applications** 

**Tested Applications:** 

**Cited Applications:** 

Species Specificity:

**Cited Species:** 

human, mouse

Positive Controls:

WB: SH-SY5Y cells, human brain tissue, mouse brain

tissue, rat brain tissue

IP: mouse brain tissue,

### **Background Information**

Rac3 is a GTPase that belongs to the RAS superfamily of small GTP-binding proteins. Members of this superfamily appear to regulate a diverse array of cellular events, including the control of cell growth, cytoskeletal reorganization, and the activation of protein kinases. There are three Rac genes in mammals, which share over 90% protein identity but differ for their pattern of expression. Rac1 is ubiquitous, Rac2 is specific for hematopoietic cells, while Rac3 is prevalent in the developing nervous system. This antibody can recognize Rac1, Rac2, and Rac3.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Brian J Rosenberg	28356423	Mol Biol Cell	WB
Erika Di Biase	32198666	Glycoconj J	WB
Sira A Franchi	29358905	Front Cell Neurosci	WB

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 16117-1-AP (RAC 1/2/3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP Result of anti-RAC 1/2/3 (IP:16117-1-AP, 3ug; Detection:16117-1-AP 1:500) with mouse brain tissue lysate 2640ug.