For Research Use Only

PGAM1 Polyclonal antibody Catalog Number: 16126-1-AP Featured Product 34

Featured Product 34 Publications



Basic Information	Catalog Number: 16126-1-AP	GenBank Accession Number: BC011678	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 300 ug/ml by	5223	WB 1:2000-1:12000	
	Nanodrop;	UNIPROT ID:	IHC 1:600-1:2400	
	Source:	P18669	IF/ICC 1:200-1:800	
	Rabbit	Full Name:		
	Isotype:	phosphoglycerate mutase 1 (brair	ו)	
	IgG Immunogen Catalog Number: AG9110	Calculated MW: 254 aa, 29 kDa		
		Observed MW: 29 kDa		
Applications	Tested Applications:	Positive C	Positive Controls:	
	WB, IHC, IF/ICC, ELISA	WB: A549 cells, HEK-293 cells, NIH/3T3 cells, Raji		
	Cited Applications:	cells, HeLa cells, HEK-293T cells, Jurkat cells, CHO		
	WB, IHC, IF, IP, CoIP	cells		
	Species Specificity: human, mouse, rat	IHC : huma breast can	in normal colon, human brain tissue, human	
	Cited Species:			
	human, mouse	IF/ICC : Ag	549 cells,	
	TE buffer pH 9.0; (*) Alternati			
	retrieval may be performed w buffer pH 6.0			
Background Information	retrieval may be performed w buffer pH 6.0 PGAM1(phosphoglycerate mutase 1) family. Phosphoglycerate mutase is reaction of 3-phosphoglycerate (3-PC PGAM1 is expressed mainly in liver, breast carcinoma, colorectal cancer, squamous cell carcinomas and also a	is also named as PGAMA,PGAM-B a widely distributed in mammalian t iA) to 2-phosphoglycerate (2-PGA) i kidney, brain and overexpressed in ung cancer, prostate cancer, oral sq ussociated with certain virus infecti- btential therapeutic target for hepat	on. PGAM1 could be developed as a useful ocellular carcinoma (PMID:20403181). This	
	retrieval may be performed w buffer pH 6.0 PGAM1(phosphoglycerate mutase 1) family. Phosphoglycerate mutase is reaction of 3-phosphoglycerate (3-PC PGAM1 is expressed mainly in liver, breast carcinoma, colorectal cancer, I squamous cell carcinomas and also a diagnostic biomarker, as well as a po antibody may also recognize PGAM2	is also named as PGAMA,PGAM-B a widely distributed in mammalian t iA) to 2-phosphoglycerate (2-PGA) i kidney, brain and overexpressed in ung cancer, prostate cancer, oral sq ussociated with certain virus infecti- btential therapeutic target for hepat	issues where it catalyzes the reversible n the glycolytic pathway. The homodimer a variety of human cancers, including uamous cell carcinoma, esophageal on. PGAM1 could be developed as a useful cocellular carcinoma (PMID:20403181). This	
	retrieval may be performed w buffer pH 6.0 PGAM1(phosphoglycerate mutase 1) family. Phosphoglycerate mutase is reaction of 3-phosphoglycerate (3-PC PGAM1 is expressed mainly in liver, breast carcinoma, colorectal cancer, I squamous cell carcinomas and also a diagnostic biomarker, as well as a po antibody may also recognize PGAM2 Author Put	is also named as PGAMA,PGAM-B a widely distributed in mammalian t iA) to 2-phosphoglycerate (2-PGA) i kidney, brain and overexpressed in ung cancer, prostate cancer, oral sq ssociated with certain virus infecti- btential therapeutic target for hepat and PGAM4 due to the high homolo	issues where it catalyzes the reversible n the glycolytic pathway. The homodimer a variety of human cancers, including uamous cell carcinoma, esophageal on. PGAM1 could be developed as a useful cocellular carcinoma (PMID:20403181). This ogy.	
Background Information	retrieval may be performed w buffer pH 6.0PGAM1(phosphoglycerate mutase 1) family. Phosphoglycerate mutase is reaction of 3-phosphoglycerate (3-PC PGAM1 is expressed mainly in liver, breast carcinoma, colorectal cancer, I squamous cell carcinomas and also a diagnostic biomarker, as well as a p antibody may also recognize PGAM2AuthorPut Rongkun Li348	is also named as PGAMA,PGAM-B a widely distributed in mammalian t A) to 2-phosphoglycerate (2-PGA) i kidney, brain and overexpressed in ung cancer, prostate cancer, oral sq ossociated with certain virus infecti- tential therapeutic target for hepat and PGAM4 due to the high homolo med ID Journal	issues where it catalyzes the reversible n the glycolytic pathway. The homodimer a variety of human cancers, including uamous cell carcinoma, esophageal on. PGAM1 could be developed as a useful cocellular carcinoma (PMID:20403181). This pgy.	
	retrieval may be performed w buffer pH 6.0 PGAM1(phosphoglycerate mutase 1) family. Phosphoglycerate mutase is reaction of 3-phosphoglycerate (3-PC PGAM1 is expressed mainly in liver, breast carcinoma, colorectal cancer, I squamous cell carcinomas and also a diagnostic biomarker, as well as a po antibody may also recognize PGAM2 Author Put Rongkun Li 348 Longzhu Ke 363	is also named as PGAMA,PGAM-B a widely distributed in mammalian t is also to 2-phosphoglycerate (2-PGA) i kidney, brain and overexpressed in ung cancer, prostate cancer, oral sq issociated with certain virus infecti- tential therapeutic target for hepat and PGAM4 due to the high homolo med ID Journal 36938 Cell Death Dis	issues where it catalyzes the reversible n the glycolytic pathway. The homodimer a variety of human cancers, including uamous cell carcinoma, esophageal on. PGAM1 could be developed as a useful cocellular carcinoma (PMID:20403181). This pgy. Application WB	
	retrieval may be performed w buffer pH 6.0 PGAM1(phosphoglycerate mutase 1) family. Phosphoglycerate mutase is reaction of 3-phosphoglycerate (3-PC PGAM1 is expressed mainly in liver, breast carcinoma, colorectal cancer, 1 squamous cell carcinomas and also a diagnostic biomarker, as well as a po antibody may also recognize PGAM2 Author Put Rongkun Li 348 Longzhu Ke 363	is also named as PGAMA,PGAM-B a widely distributed in mammalian t is also 2-phosphoglycerate (2-PGA) i kidney, brain and overexpressed in ung cancer, prostate cancer, oral sq issociated with certain virus infecti- tential therapeutic target for hepat and PGAM4 due to the high homolo med ID Journal 36938 Cell Death Dis 35636 Cell Biol Int 02531 Bioengineered er shipment.	issues where it catalyzes the reversible n the glycolytic pathway. The homodimer a variety of human cancers, including uamous cell carcinoma, esophageal on. PGAM1 could be developed as a useful cocellular carcinoma (PMID:20403181). This pgy. Application WB WB	


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com

in USA), or 1(312) 455-8498 (outside USA)

W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16126-1-AP (PGAM1 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 16126-1-AP (PGAM1 antibody) at dilution of 1:1200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 16126-1-AP (PGAM1 antibody) at dilution of 1:1200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using PGAM1 antibody (16126-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).