For Research Use Only

ACSS3 Polyclonal antibody

Catalog Number:16204-1-AP







Basic Information	Catalog Number: 16204-1-AP	GenBank Accession Number: BC009317	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 500 ug/ml by		WB 1:1000-1:5000	
	Nanodrop;	UNIPROT ID:	IHC 1:20-1:200	
	Source:	Q9H6R3	IF/ICC 1:200-1:800	
	Rabbit	Full Name:		
	Isotype: IgG	acyl-CoA synthetase short-chain family member 3		
	Immunogen Catalog Number:	Calculated MW:		
	AG9173	686 aa, 75 kDa		
		Observed MW: 70-75 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, IF/ICC, ELISA	WB: HepG	WB : HepG2 cells, mouse kidney tissue, rat kidney	
	Cited Applications:	tissue		
	WB, IHC	IHC : huma	n liver tissue,	
	Species Specificity: human, mouse, rat	IF/ICC : U2	IF/ICC : U2OS cells,	
	Cited Species: human, mouse			
	Note-IHC: suggested antigen i TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0	vely, antigen		
Background Information	binding enzyme family. It activates	acetate so that it can be used for lip tains 4 of 5 motifs characteristic of a	cyl-CoA synthetases. This protein has 2	
	binding enzyme family. It activates deduced 686-amino acid protein con isoforms produced by alternative spl	acetate so that it can be used for lip tains 4 of 5 motifs characteristic of a	d synthesis or for energy generation. Th acyl-CoA synthetases. This protein has 2	
	binding enzyme family. It activates deduced 686-amino acid protein con isoforms produced by alternative spl Author Pu	acetate so that it can be used for lip tains 4 of 5 motifs characteristic of a icing. The full length protein has a t	d synthesis or for energy generation. Th acyl-CoA synthetases. This protein has 2 ransit peptide with 29 amino acids.	
Background Information	binding enzyme family. It activates deduced 686-amino acid protein con isoforms produced by alternative spl Author Pu Karolin Weitkunat 34	acetate so that it can be used for lipi tains 4 of 5 motifs characteristic of a icing. The full length protein has a t bmed ID Journal	d synthesis or for energy generation. Th acyl-CoA synthetases. This protein has 2 ransit peptide with 29 amino acids. Application	
	binding enzyme family. It activates deduced 686-amino acid protein con isoforms produced by alternative spl Author Pu Karolin Weitkunat 34 Dongsheng Shen 39	acetate so that it can be used for lip tains 4 of 5 motifs characteristic of a icing. The full length protein has a t bmed ID Journal .064336 Nutrients	d synthesis or for energy generation. Th acyl-CoA synthetases. This protein has 2 ransit peptide with 29 amino acids. Application WB	
	binding enzyme family. It activates deduced 686-amino acid protein con isoforms produced by alternative spl Author Pu Karolin Weitkunat 34 Dongsheng Shen 39	acetate so that it can be used for lipitains 4 of 5 motifs characteristic of a icing. The full length protein has a tomed ID Journal 064336 Nutrients 754905 Cancer Genet 146936 Cell Metab ter shipment.	d synthesis or for energy generation. Th acyl-CoA synthetases. This protein has 2 ransit peptide with 29 amino acids. Application WB WB	
Notable Publications	binding enzyme family. It activates deduced 686-amino acid protein con isoforms produced by alternative spl Author Pu Karolin Weitkunat 34 Dongsheng Shen 39 Bing Zhou 39 Storage: Storage Storage Storage Buffer:	acetate so that it can be used for lipitains 4 of 5 motifs characteristic of a icing. The full length protein has a tomed ID Journal 064336 Nutrients 754905 Cancer Genet 146936 Cell Metab ter shipment.	d synthesis or for energy generation. Th acyl-CoA synthetases. This protein has 2 ransit peptide with 29 amino acids. Application WB WB	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data







Various lysates were subjected to SDS PAGE followed by western blot with 16204-1-AP (ACSS3 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human liver using 16204-1-AP (ACSS3 antibody) at dilution of 1:100 (under 10x lens).


Immunohistochemical analysis of paraffinembedded human liver using 16204-1-AP (ACSS3 antibody) at dilution of 1:100 (under 40x lens).



WB result of ACSS3 antibody (16204-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ACSS3 transfected HepG2 cells.



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using ACSS3 antibody (16204-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using ACSS3 antibody (16204-1-AP) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).