

For Research Use Only

TBRG4 Polyclonal antibody

Catalog Number: 16245-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number: 16245-1-AP	GenBank Accession Number: BC017235	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 400 µg/ml by Nanodrop and 327 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 9238	Recommended Dilutions: WB 1:500-1:3000 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
Source: Rabbit	Full Name: transforming growth factor beta regulator 4	IHC 1:100-1:400
Isotype: IgG	Calculated MW: 631 aa, 71 kDa	
Immunogen Catalog Number: AG9176	Observed MW: 60 kDa, 71 kDa	

Applications

Tested Applications:
IF, IHC, IP, WB, ELISA

Cited Applications:
HITS-CLIP, WB

Species Specificity:
human, mouse, rat

Cited Species:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HepG2 cells, HeLa cells, human liver tissue, mouse ovary tissue

IP : HepG2 cells,

IHC : human liver cancer tissue,

Background Information

TBRG4 is also named as CPR2(Cell cycle progression restoration protein 2), FASTKD4(FAST kinase domain-containing protein 4), KIAA0948 and belongs to the FAST kinase family. This protein contains multiple putative leucine zipper domains characteristic of transcription factors and it may play a role in cell cycle progression(PMID:9383053). RT-PCR ELISA detected moderate expression in ovary, and little to no expression in all other tissues and specific brain regions examined(PMID:10231032).TBRG4 has an mature form with MW of 60 kDa produced by transit peptide missing.

Notable Publications

Author	Pubmed ID	Journal	Application
Akira Ohkubo	34748562	PLoS Genet	WB,HITS-CLIP
Claudio Asencio	30035255	Life Sci Alliance	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 16245-1-AP (TBRG4 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




IP Result of anti-TBRG4 (IP:16245-1-AP, 4ug; Detection:16245-1-AP 1:500) with HepG2 cells lysate 1600ug.



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 16245-1-AP (TBRG4 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 16245-1-AP (TBRG4 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



U2OS cells, fixed with 4% pF/PBS and post-fixed with -20 methanol, and stained for TIAR (green), anti-TBRG4 (red, cat# 16245-1-AP), and human anti-mitochondria autoantiserum (blue). Note that the TBRG4 signal is strongly mitochondrial, but also nucleolar.

IF result of anti-TBRG4 (16245-1-AP) with U2OS cells.