

For Research Use Only

# ORM1 Polyclonal antibody

Catalog Number: 16439-1-AP

Featured Product

14 Publications



## Basic Information

### Catalog Number:

16439-1-AP

### Size:

150ul, Concentration: 900 ug/ml by Nanodrop and 447 ug/ml by Bradford method using BSA as the standard;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG9758

### GenBank Accession Number:

BC026238

### GeneID (NCBI):

5004

### UNIPROT ID:

P02763

### Full Name:

orosomucoid 1

### Calculated MW:

201 aa, 24 kDa

### Observed MW:

40-47 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

### Cited Applications:

WB, IHC, IF, CoIP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

WB : mouse liver tissue, mouse bladder tissue

IP : human plasma tissue,

IHC : human liver tissue, human normal colon, human liver cancer tissue

IF/ICC : HepG2 cells,

## Background Information

Alpha-1-acid glycoprotein 1 (AGP1), also called orosomucoid-1 (ORM1), is a glycoprotein synthesized mostly by hepatocytes and present in human plasma. ORM1 is an acute-phase reactant protein controlled by glucocorticoids, interleukin-1 and interleukin-6, and increase up to 5-50 times upon infection and/or inflammation. Anti-apoptotic effect and role as immunomodulator of ORM have been reported. ORM is an important carrier for synthetic drugs and influences their distribution and availability in the body. This antibody recognizes a band about 44 kDa in human plasma which may be due to the glycosylation of ORM1 or the dimer formation of the protein.

## Notable Publications

Author	Pubmed ID	Journal	Application
Mehrpouya B Mobin	27665711	Nat Commun	WB
Bing Zhou	36050503	Nat Metab	WB
Luo Qiong	34654351	Bioengineered	IHC

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




mouse liver tissue were subjected to SDS PAGE followed by western blot with 16439-1-AP (ORM1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 16439-1-AP (ORM1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 16439-1-AP (ORM1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-ORM1 (IP:16439-1-AP, 3ug; Detection:16439-1-AP 1:300) with human plasma lysate 100ug.



$1 \times 10^6$  HepG2 cells were intracellularly stained with 0.4 ug ORM1 Polyclonal antibody (16439-1-AP) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.4 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 16439-1-AP (ORM1 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using ORM1 antibody (16439-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).