

For Research Use Only

# OCIAD1 Polyclonal antibody

Catalog Number: 16634-1-AP

2 Publications



## Basic Information

### Catalog Number:

16634-1-AP

### Size:

150ul, Concentration: 350 ug/ml by Nanodrop and 313 ug/ml by Bradford method using BSA as the standard;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG9977

### GenBank Accession Number:

BC003409

### GeneID (NCBI):

54940

### UNIPROT ID:

Q9NX40

### Full Name:

OCIA domain containing 1

### Calculated MW:

28 kDa

### Observed MW:

29-35 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

### Cited Applications:

WB, IHC, IF

### Species Specificity:

human

### Cited Species:

human

### Positive Controls:

WB : HeLa cells, MCF-7 cells, human placenta tissue

IP : HeLa cells,

IHC : human liver cancer tissue, human kidney tissue, human pancreas cancer tissue, human thyroid cancer tissue

IF/ICC : HepG2 cells, HeLa cells

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

OCIAD1 was first identified by immunoscreening of an ovarian carcinoma cDNA expression library with ascites fluid from ovarian cancer patients (PMID: 11162530). OCIAD1 has been reported as a key player in ovarian cancer cell adhesion, as well as a key player in generating ovarian cancer recurrence (PMID: 18328549; 20515946). In addition to its roles in cancer, OCIAD1 participates in maintaining stem cell potency by regulating the Jak/STAT pathway (PMID: 23972987). Several alternatively spliced forms of OCIAD1 gene have been identified. The longest form (1.4 kb) is predicted to encode for a 27.6 kDa protein of 245 amino acids. This antibody detects OCIAD1 with an apparent molecular weight of ~35 kDa as has been demonstrated by several researches (PMID: 27345969; 27345976).

## Notable Publications

Author	Pubmed ID	Journal	Application
Huong T L Tran	32697788	PLoS One	WB,IF
Nagata Chigusa C	22726067	Pathol Int	IHC

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16634-1-AP (OCIAD1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 16634-1-AP (OCIAD1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 16634-1-AP (OCIAD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using OCIAD1 antibody (16634-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).



IP result of anti-OCIAD1 (IP:16634-1-AP, 4ug; Detection:16634-1-AP 1:300) with HeLa cells lysate 3200ug.