

For Research Use Only

# MRPL10 Polyclonal antibody

Catalog Number: 16652-1-AP **1 Publications**



## Basic Information

<b>Catalog Number:</b> 16652-1-AP	<b>GenBank Accession Number:</b> BC052601	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul, Concentration: 400 µg/ml by Nanodrop and 207 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 124995	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IP 0.5-4.0 ug for IP and 1:200-1:1000 for WB
<b>Source:</b> Rabbit	<b>Full Name:</b> mitochondrial ribosomal protein L10	<b>IHC 1:20-1:200</b>
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 261 aa, 29 kDa	<b>IF 1:50-1:500</b>
<b>Immunogen Catalog Number:</b> AG10087	<b>Observed MW:</b> 29 kDa	

## Applications

<b>Tested Applications:</b> IF, IHC, IP, WB, ELISA	<b>Positive Controls:</b> WB : HEK-293 cells, HEK-293 mitochondrial
<b>Cited Applications:</b> WB	<b>IP :</b> mouse liver tissue,
<b>Species Specificity:</b> human, mouse, rat	<b>IHC :</b> human testis tissue,
<b>Cited Species:</b> human	<b>IF :</b> HepG2 cells,

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

### Notable Publications

Author	Pubmed ID	Journal	Application
Luis Daniel Cruz-Zaragoza	34672953	Cell	WB

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16652-1-AP (MRPL10 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human testis using 16652-1-AP (MRPL10 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human testis using 16652-1-AP (MRPL10 antibody) at dilution of 1:50 (under 40x lens).



IP result of anti-MRPL10 (IP:16652-1-AP, 4ug; Detection:16652-1-AP 1:300) with mouse liver tissue lysate 6000 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using MRPL10 antibody (16652-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).