

CD3 Delta Polyclonal antibody

Catalog Number: 16669-1-AP

6 Publications

Basic Information

Catalog Number:

16669-1-AP

Size:

150ul , Concentration: 300 µg/ml by Nanodrop and 260 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG10148

GenBank Accession Number:

BC070321

GeneID (NCBI):

915

Full Name:

CD3d molecule, delta (CD3-TCR complex)

Calculated MW:

171 aa, 19 kDa

Observed MW:

20-25 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:4000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB

IF 1:50-1:500

Applications

Tested Applications:

FC, IF, IP, WB, ELISA

Cited Applications:

IF, IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB : Jurkat cells, mouse thymus tissue, rat thymus tissue, MOLT-4 cells

IP : Jurkat cells,

IF : human tonsillitis tissue,

Background Information

CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuji Fukushima	36130493	Cell Rep	WB,IP
Marie Klintman	27179111	Clin Cancer Res	
Binghao Li	29735547	Cancer Res	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16669-1-AP (CD3 Delta antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




1X10⁶ Jurkat cells were stained with 0.5ug CD3 delta antibody (16669-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD3 Delta antibody (16669-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD3 Delta antibody (16669-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP Result of anti-CD3 delta (IP:16669-1-AP, 4ug; Detection:16669-1-AP 1:800) with Jurkat cells lysate 2800ug.