

For Research Use Only

NAA40 Polyclonal antibody

Catalog Number: 16698-1-AP

1 Publications



Basic Information

Catalog Number: 16698-1-AP	GenBank Accession Number: BC052298	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 450 ug/ml by Nanodrop;	GeneID (NCBI): 79829	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	UNIPROT ID: Q86UY6	IHC 1:50-1:500
Isotype: IgG	Full Name: N-acetyltransferase 11 (GCN5-related, putative)	IF/ICC 1:50-1:500
Immunogen Catalog Number: AG10092	Calculated MW: 237 aa, 27 kDa	
	Observed MW: 27 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA	Positive Controls: WB : HEK-293 cells, HeLa cells, HepG2 cells, mouse spleen tissue
Cited Applications: WB, IF	IHC : human ovarian cancer,
Species Specificity: human, mouse, rat	IF/ICC : U2OS cells,
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

N-alpha-acetyltransferase 40 (NAA40, also known as NatD, Nat4, or Patt1) is a highly specific epigenetic enzyme catalyzing the transfer of an acetyl moiety at the N-terminal end of histones H4 and H2A (PMID: 26666750). NAA40 is significantly upregulated in primary CRC tissues and promotes CRC cell growth both in vitro and in xenograft tumor models (PMID: 30858358). Western blot analysis detected NAA40 at an apparent molecular mass of 27 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Chunfei Li	39757327	Cell Biol Toxicol	WB,IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16698-1-AP (NAT11 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human ovarian cancer slide using 16698-1-AP (NAA40 antibody) at dilution of 1:100 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using NAA40 antibody (16698-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).