

For Research Use Only

GSTA3 Polyclonal antibody

Catalog Number: 16703-1-AP

5 Publications



Basic Information

Catalog Number:

16703-1-AP

Size:

150ul, Concentration: 600 µg/ml by Nanodrop and 307 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG10101

GenBank Accession Number:

BC020619

GeneID (NCBI):

2940

Full Name:

glutathione S-transferase alpha 3

Calculated MW:

222 aa, 25 kDa

Observed MW:

25 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IP 0.5-4.0 µg for IP and 1:500-1:1000 for WB

IF 1:10-1:100

Applications

Tested Applications:

IF, IP, WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

mouse, rat

Positive Controls:

WB : mouse testis tissue, rat testis tissue, human testis tissue

IP : mouse testis tissue,

IF : MCF-7 cells,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Yoko Ito	33237957	PLoS One	WB
LaLida Rojanathammanee	25001375	Age (Dordr)	WB
Li Qin Zhang	29684358	Am J Pathol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Immunofluorescent analysis of MCF-7 cells, using GSTA3 antibody 16703-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



IP Result of anti-GSTA3 (IP:16703-1-AP, 4ug; Detection:16703-1-AP 1:500) with mouse testis tissue lysate 4800ug.



mouse testis tissue was subjected to SDS PAGE followed by western blot with 16703-1-AP (GSTA3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.