

For Research Use Only

SPOP Polyclonal antibody

Catalog Number: 16750-1-AP

Featured Product

70 Publications



Basic Information

Catalog Number: 16750-1-AP	GenBank Accession Number: BC003385	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 µg/ml by Nanodrop;	GeneID (NCBI): 8405	Recommended Dilutions: WB 1:5000-1:50000 IF 1:50-1:500
Source: Rabbit	UNIPROT ID: O43791	
Isotype: IgG	Full Name: speckle-type POZ protein	
Immunogen Catalog Number: AG10215	Calculated MW: 374 aa, 42 kDa	
	Observed MW: 42 kDa	

Applications

Tested Applications: WB, IF, ELISA	Positive Controls: WB : HeLa cells, HepG2 cells, PC-3 cells IF : HepG2 cells,
Cited Applications: WB, IP, IF, RIP, IHC, CoIP	
Species Specificity: human, mouse, rat	
Cited Species: human, rat, mouse	

Background Information

The SPOP (TEF2) protein was previously identified as an autoantigen in a patient with scleroderma pigmentosum. SPOP (speckle-type POZ protein), also known as TEF2, HIB homolog 1 or Roadkill homolog 1, is a member of the Tdpoz family containing one N-terminal MATH (Mepirin and TRAF Homology) domain and one C-terminal BTB/POZ domain. SPOP can exist as a homodimer and is expressed in a variety of tissues localizing to the nucleus. BTB-mediated SPOP dimers form linear oligomers via BACK domain dimerization, and we determine the concentration-dependent populations of the resulting oligomeric species (PMID: 27220849). Through an interaction with CUL-3, SPOP is involved in ubiquitinylation and protein degradation. SPOP specifically interacts with CUL-3 via its BTB/POZ domain and recruits substrates to the CUL-3-based ubiquitin ligase via its MATH domain. Substrates recruited by SPOP and targeted for ubiquitylation via the CUL-3/SPOP complex include PDX-1, Bmi-1, MacroH2A, PIPK II β and Daxx. These substrates are subsequently degraded by the proteasome. In addition, SPOP itself becomes ubiquitylated by the CUL-3-based ubiquitin ligase and is targeted for proteasomal degradation.

Notable Publications

Author	Pubmed ID	Journal	Application
Carley Snoznik	34593637	Proc Natl Acad Sci U S A	WB
Jianong Zhang	34588438	Nat Commun	WB
Lan Zhang	34586738	Clin Transl Med	WB,IHC,RIP

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 16750-1-AP (SPOP antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 16750-1-AP (SPOP antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 16750-1-AP (SPOP antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.