

For Research Use Only

Vitamin D binding protein Polyclonal antibody



Catalog Number: 16922-1-AP

8 Publications

Basic Information

Catalog Number: 16922-1-AP	GenBank Accession Number: BC057228	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 300 µg/ml by Nanodrop and 200 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2638	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500
Source: Rabbit	Full Name: group-specific component (vitamin D binding protein)	
Isotype: IgG	Calculated MW: 474 aa, 53 kDa	
Immunogen Catalog Number: AG10286	Observed MW: 52-58 kDa	

Applications

Tested Applications:
IHC, WB, ELISA

Cited Applications:
IF, IHC, WB

Species Specificity:
human, mouse, rat

Cited Species:
human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human blood tissue, human blood, rat eye tissue, mouse eye tissue, PC-3 cells, human plasma, mouse testis tissue

IHC: human liver tissue,

Background Information

Vitamin D binding protein is a sparsely glycosylated serum protein responsible for highly specific binding and tissue-specific delivery of vitamin D and its metabolites. In addition, it is also an actin scavenger, and is the precursor to the immunomodulatory protein, Gc-MAF. Vitamin D binding protein has been proposed to have significant roles in C5a chemotaxis, osteoclast development and possibly in macrophage activation/recruitment.

Notable Publications

Author	Pubmed ID	Journal	Application
Manish Kumar Yadav	36284768	Mol Ther Methods Clin Dev	WB
Yang Lu	26791873	Proteomics	
Yichen Guo	28656274	Mol Med Rep	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human blood were subjected to SDS PAGE followed by western blot with 16922-1-AP (vitamin D binding protein antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 16922-1-AP (Vitamin D binding protein antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 16922-1-AP (Vitamin D binding protein antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



rat eye tissue were subjected to SDS PAGE followed by western blot with 16922-1-AP (Vitamin D binding protein antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.