For Research Use Only

## AP1B1 Polyclonal antibody Catalog Number: 16932-1-AP 5 Publications





Basic Information	Catalog Number: 16932-1-AP	GenBank Accession Number: BC046242		Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:2400 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate	
	Size: GenelD (NCBI):				
	150ul , Concentration: 750 ug/ml by	162			
	Nanodrop and 333 ug/ml by Bradford method using BSA as the standard;				
	Source:	Full Name:		IF/ICC 1:50-1:500	
	Rabbit	adaptor-related protein complex 1,			
	Isotype:	beta 1 subunit			
	IgG Immunogen Catalog Number: AG10417	Calculated MW:			
		919 aa, 101 kDa			
		Observed MW: 101 kDa			
Applications	Tested Applications:	Positive Controls:			
	Cited Applications: IP : HFK-20		WB: HEK-293	cells,	
			IP : HEK-293 c	ells,	
	WB		IF/ICC : HepG2 cells,		
	Species Specificity: human				
	Cited Species:				
	human				
	nanan				
Background Information	AP1B1 is a subunit of the adaptor con complexes are cytosolic heterotetran	hers that mediate the so the cytoplasmic face of hrin to the membrane a	orting of memb coated vesicle	s located at the Golgi complex, where	
	AP1B1 is a subunit of the adaptor com complexes are cytosolic heterotetran endocytic pathways. AP-1 is found at mediates both the recruitment of clat cytosolic tails of transmembrane rece	hers that mediate the so the cytoplasmic face of hrin to the membrane a	orting of memb coated vesicle nd the recogni	rane proteins in the secretory and s located at the Golgi complex, where	
	AP1B1 is a subunit of the adaptor com complexes are cytosolic heterotetran endocytic pathways. AP-1 is found at mediates both the recruitment of clat cytosolic tails of transmembrane reco Author Pu	ners that mediate the so the cytoplasmic face of hrin to the membrane a optors.	orting of memb coated vesicle nd the recogni	rane proteins in the secretory and s located at the Golgi complex, where tion of sorting signals within the	
Background Information	AP1B1 is a subunit of the adaptor com complexes are cytosolic heterotetran endocytic pathways. AP-1 is found at mediates both the recruitment of clat cytosolic tails of transmembrane rece Author Pu Arunas Kvainickas 28	hers that mediate the so the cytoplasmic face of hrin to the membrane a sptors. bmed ID Journ 935632 J Cell	orting of memb coated vesicle nd the recogni	rane proteins in the secretory and s located at the Golgi complex, where tion of sorting signals within the Application	
	AP1B1 is a subunit of the adaptor com complexes are cytosolic heterotetran endocytic pathways. AP-1 is found at mediates both the recruitment of clat cytosolic tails of transmembrane reco Author Pu Arunas Kvainickas 28 Lynn M Boyden 31	hers that mediate the so the cytoplasmic face of hrin to the membrane a sptors. bmed ID Journ 935632 J Cell	orting of memb coated vesicle nd the recogni al Biol Hum Genet	rane proteins in the secretory and s located at the Golgi complex, where tion of sorting signals within the Application WB	
	AP1B1 is a subunit of the adaptor com complexes are cytosolic heterotetran endocytic pathways. AP-1 is found at mediates both the recruitment of clat cytosolic tails of transmembrane reco Author Pu Arunas Kvainickas 28 Lynn M Boyden 31	hers that mediate the so the cytoplasmic face of hrin to the membrane a sptors. bmed ID Journ 935632 J Cell 630788 Am J	orting of memb coated vesicle nd the recogni al Biol Hum Genet	rane proteins in the secretory and s located at the Golgi complex, where tion of sorting signals within the Application WB WB	
Notable Publications	AP1B1 is a subunit of the adaptor com complexes are cytosolic heterotetran endocytic pathways. AP-1 is found at mediates both the recruitment of clat cytosolic tails of transmembrane reco Author Pu Arunas Kvainickas 28 Lynn M Boyden 31	rers that mediate the so the cytoplasmic face of hrin to the membrane a sptors. bmed ID Journ 935632 J Cell 630788 Am J 879413 Nat G er shipment.	orting of memb coated vesicle nd the recogni al Biol Hum Genet	rane proteins in the secretory and s located at the Golgi complex, where tion of sorting signals within the Application WB WB	
	AP1B1 is a subunit of the adaptor com complexes are cytosolic heterotetran endocytic pathways. AP-1 is found at mediates both the recruitment of clat cytosolic tails of transmembrane reco Author Pu Arunas Kvainickas 28 Lynn M Boyden 31 Antoine Rebendenne 35 Storage: Storage Store at -20°C. Stable for one year aft Storage Buffer:	rers that mediate the so the cytoplasmic face of hrin to the membrane a sptors. bmed ID Journ 935632 J Cell 630788 Am J 879413 Nat G er shipment. % glycerol pH 7.3.	orting of memb coated vesicle nd the recogni al Biol Hum Genet	rane proteins in the secretory and s located at the Golgi complex, where tion of sorting signals within the Application WB WB	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





HEK-293 cells were subjected to SDS PAGEIP result of anti-AP1B1 (IP:16932-1-AP, 4ug;followed by western blot with 16932-1-AP (AP1B1Detection:16932-1-AP 1:300) with HEK-293 cellsantibody) at dilution of 1:500 incubated at roomlysate 2320 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 16932-1-AP (AP1B1 antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated Goat Anti-Rabbit IgG(H+L).