For Research Use Only

# TADA2B Polyclonal antibody

Catalog Number:17367-1-AP

Featured Product

2 Publications



### **Basic Information**

Catalog Number: 17367-1-AP Size: 150ul , Concentration: 750 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG10783

GenBank Accession Number: BC013164 GeneID (NCBI): 93624 UNIPROT ID: Q86TJ2 Full Name: transcriptional adaptor 2 (ADA2 homolog, yeast)-beta Calculated MW: 420 aa, 48 kDa Observed MW:

55-62 kDa

#### Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:1000-1:4000 IHC 1:200-1:800

# Applications

Tested Applications: WB, IHC, ELISA Cited Applications: WB Species Specificity: human Cited Species: human Note-IHC: suggested antigen retrieval with

TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB : HeLa cells, HepG2 cells, K-562 cells, MCF-7 cells IHC : human ovary cancer tissue,

## Notable Publications

Author	Pubmed ID	Journal	Application
Rui Guo	32315601	Mol Cell	WB
Clare F Malone	38820154	Sci Adv	WB

### Storage


Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

# Selected Validation Data





Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 17367-1-AP (TADA2B antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Various lysates were subjected to SDS PAGE followed by western blot with 17367-1-AP (TADA2B antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.