

For Research Use Only

PADI4/PAD4 Polyclonal antibody

Catalog Number: 17373-1-AP

Featured Product

45 Publications



Basic Information

Catalog Number:

17373-1-AP

Size:

150ul, Concentration: 1000 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG11068

GenBank Accession Number:

BC025718

GeneID (NCBI):

23569

UNIPROT ID:

Q9UM07

Full Name:

peptidyl arginine deiminase, type IV

Calculated MW:

663 aa, 74 kDa

Observed MW:

67 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: SKOV-3 cells, mouse spleen tissue

IP: mouse spleen tissue,

IHC: mouse spleen tissue,

IF/ICC: A549 cells,

Background Information

PADI4, also named PADI5 and PDI5, catalyzes the citrullination/deimination of arginine residues of proteins such as histones, thereby playing a key role in histone code and regulation of stem cell maintenance. PADI4 is expressed in hematologic and rheumatoid arthritis synovial tissues. PADI4 haplotype associated with susceptibility to rheumatoid arthritis increases the production of citrullinated peptides, leading, in a cytokine-rich milieu, to a break in tolerance to citrullinated peptides processed and presented in the appropriate HLA context. Immunohistochemistry detected significant PADI4 expression in various malignancies and PADI4 expression was not detected in non-tumor tissues. (PMID: 19183436)

Notable Publications

Author	Pubmed ID	Journal	Application
Xinyan Wu	34721438	Front Immunol	WB
Szilvia Török	34884536	Int J Mol Sci	WB
Atsumu Osada	33238047	Clin Exp Immunol	IHC, IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




SKOV-3 cells were subjected to SDS PAGE followed by western blot with 17373-1-AP (PADI4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 17373-1-AP (PADI4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed A549 cells using PADI4 antibody (17373-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Anti-Human PADI4 (17373-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).




Immunohistochemical analysis of paraffin-embedded mouse spleen tissue slide using 17373-1-AP (PADI4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse spleen tissue were subjected to SDS PAGE followed by western blot with 17373-1-AP (PADI4 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



IP result of anti-PADI4 (IP:17373-1-AP, 4ug; Detection:17373-1-AP 1:500) with mouse spleen tissue lysate 4800ug.



Immunofluorescent analysis of (4% PFA) fixed A549 cells using PADI4 antibody (17373-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).