For Research Use Only

DDX4,VASA Polyclonal antibody

Catalog Number: 17545-1-AP

1 Publications



Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:20000-1:100000

IHC 1:200-1:800

Basic Information

Catalog Number: GenBank Accession Number: 17545-1-AP BC047455

Size: GeneID (NCBI):

150ul , Concentration: 900 μ g/ml by 54514 Nanodrop and 460 μ g/ml by Bradford Full Name:

method using BSA as the standard; DEAD (Asp-Glu-Ala-Asp) box

Source: polypeptide 4
Rabbit Calculated MW:

Isotype: 690aa,76 kDa; 724aa,79 kDa

IgG Observed MW:
Immunogen Catalog Number: 79 kDa

AG11647

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse testis tissue, rat testis tissue

IHC: mouse testis tissue,

Background Information

DEAD box proteins are characterized by nine conserved sequence motifs located on two functional domains. Domain I contains six of these motifs, including the Q motif and the Walker A motif, motifs Ia and Ib, the Walker B motif, and motif III, which may act to link ATPase and helicase activities of the protein [PMID:21653890]. DDX4, a member of the DEAD box family of ATP-dependent RNA helicases, plays a central role in several aspects of germ cell development. Its function is not only required during gametogenesis in the adult but is also essential for the specification of the germ cell lineage during embryogenesis [PMID:20016130].

Notable Publications

Author	Pubmed ID	Journal	Application
Larissa Zarate-Garcia	27301892	Sci Rep	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 17545-1-AP (DDX4,VASA antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



rat testis tissue were subjected to SDS PAGE followed by western blot with 17545-1-AP (DDX4,VASA antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 17545-1-AP (DDX4,VASA antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).