

For Research Use Only

# Beta Catenin Polyclonal antibody

Catalog Number: 17565-1-AP

Featured Product

121 Publications



## Basic Information

### Catalog Number:

17565-1-AP

### Size:

150ul, Concentration: 600 ug/ml by Nanodrop and 333 ug/ml by Bradford method using BSA as the standard;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG11763

### GenBank Accession Number:

BC058926

### GeneID (NCBI):

1499

### ENSEMBL Gene ID:

ENSG00000168036

### UNIPROT ID:

P35222

### Full Name:

catenin (cadherin-associated protein), beta 1, 88kDa

### Calculated MW:

781 aa, 86 kDa

### Observed MW:

92 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, ELISA

### Cited Applications:

WB, IHC, IF, IP, CoIP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB**: HEK-293 cells, NIH/3T3 cells, HeLa cells, mouse brain tissue, A431 cells, MCF-7 cells, mouse liver tissue, rat brain tissue, C6 cells, rat liver tissue

**IP**: HEK-293 cells,

**IHC**: human liver cancer tissue, human breast cancer tissue

**IF/ICC**: MCF-7 cells, T-47D cells, U-251 cells

## Background Information

$\beta$ -Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein.  $\beta$ -Catenin was originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to  $\alpha$ -catenin and the actin cytoskeleton. Besides its essential role in the AJs,  $\beta$ -catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/ $\beta$ -catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of  $\beta$ -catenin activity is associated with multiple diseases including cancers. (PMID: 22617422; 18334222)

## Notable Publications

Author	Pubmed ID	Journal	Application
Hai-Yan Wang	34588618	Acta Pharmacol Sin	WB
Jiao Wu	34517345	Aging (Albany NY)	WB
Tianfu Li	36089118	Cancer Lett	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 17565-1-AP (Beta Catenin antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 17565-1-AP (beta-Catenin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 17565-1-AP (beta-Catenin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-Beta Catenin (IP:17565-1-AP, 4ug; Detection:17565-1-AP 1:5000) with HEK-293 cells lysate 1920 ug.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using Beta Catenin antibody (17565-1-AP) at dilution of 1:400 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L), Alpha Tubulin antibody (66031-1-Ig, Clone: 1E4C11, green).



1X10<sup>6</sup> MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human Beta Catenin (17565-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).