#### For Research Use Only

# ST7L Polyclonal antibody

Catalog Number: 17567-1-AP 4 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Antigen affinity purification

17567-1-AP

Size:

BC065501 GeneID (NCBI):

Recommended Dilutions:

150ul , Concentration: 240 ug/ml by

54879

WB 1:500-1:3000

Nanodrop and 233 ug/ml by Bradford  $\,$  UNIPROT ID:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

method using BSA as the standard;

Q8TDW4

protein lysate

Source:

Full Name:

Rabbit suppression of tumorigenicity 7 like Isotype:

Immunogen Catalog Number:

Calculated MW: 575 aa, 65 kDa

AG11768

Observed MW:

60-70 kDa

**Applications** 

**Tested Applications:** 

WB, IP, ELISA

Cited Applications:

WB, IHC, CoIP

Species Specificity:

mouse, human

**Cited Species:** 

human

Positive Controls:

WB: LO2 cells, MCF-7 cells, T-47D cells

IP: mouse testis tissue.

## **Background Information**

ST7 (suppressor of tumorigenicity 7) gene has been proposed to be a tumor suppressor gene, product of which is a multi-pass membrane protein belonging to the ST7 family. ST7L (suppressor of tumorigenicity 7 protein-like) gene encodes a multi-pass membrane protein also belonging to the ST7 family, 8 isoforms produced by alternative splicing have been described. However, the molecular function of this gene product has not been determined so far.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Shaoying Yan	30176933	J Exp Clin Cancer Res	WB
Likun Zhuang	28518144	Cell Death Dis	WB,CoIP
Shengjie Li	28902356	Int J Mol Med	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

### Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17567-1-AP (ST7L antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



IP result of anti-ST7L (IP:17567-1-AP, 3ug; Detection:17567-1-AP 1:1500) with mouse testis tissue lysate 5000ug.