For Research Use Only

JNK Polyclonal antibody

Catalog Number: 17572-1-AP

Featured Product

29 Publications



Basic Information

Catalog Number: GenBank Accession Number:

17572-1-AP BC065516
Size: GeneID (NCBI):

150ul , Concentration: 450 ug/ml by 5602
Nanodrop; UNIPROT ID:
Source: P53779

Rabbit Full Name:
Isotype: mitogen-activated protein kinase 10

IgG Calculated MW:
Immunogen Catalog Number: 464 aa, 53 kDa
AG11785 Observed MW:

45-48 kDa, 54-57 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Cited Applications: WB, IHC, IF Species Specificity: human, mouse, rat

Cited Species: human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Antigen affinity purification

Recommended Dilutions: WB 1:1000-1:8000 IHC 1:50-1:500 IF/ICC 1:200-1:800

Positive Controls:

WB: mouse brain tissue, mouse cerebellum tissue, rat

brain tissue

IHC: human liver tissue, IF/ICC: HepG2 cells,

Background Information

JNK, also named MAPK10 (mitogen-activated protein kinase 10), belongs to MAP kinase family. MAP kinases act as integration points for multiple biochemical signals, and thus are involved in a wide variety of cellular processes, such as proliferation, differentiation, transcription regulation and development. This kinase is mainly expressed in a subset of neurons in the nervous system, and is activated by threonine and tyrosine phosphorylation. JNK has 3 isoforms of 48-53 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Yuan-Jian Song	30412737	Behav Brain Res	WB
Xiaochun Yang	25044439	Anat Rec (Hoboken)	WB
Ya-Sai Sun	32092821	Biomed Pharmacother	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 17572-1-AP (MAPK10 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using JNK antibody (17572-1-AP) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 17572-1-AP (MAPK10 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 17572-1-AP (MAPK10 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).