### For Research Use Only

# SLC39A9 Polyclonal antibody

Catalog Number: 17607-1-AP



**Purification Method:** 

WB 1:500-1:2000 IHC 1:50-1:500

Antigen affinity purification

Recommended Dilutions:

**Basic Information** 

Catalog Number: GenBank Accession Number:

17607-1-AP BC047682 GeneID (NCBI): 150ul , Concentration: 400 µg/ml by 55334

Source: solute carrier family 39 (zinc Rabbit transporter), member 9 Isotype: Calculated MW: IgG 307 aa, 32 kDa

Immunogen Catalog Number: Observed MW: 25 kDa AG11752

**Applications** 

**Tested Applications:** Positive Controls:

IHC, WB, ELISA WB: HUVEC cells, Species Specificity:

IHC: human placenta tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

### **Background Information**

Zinc transporter ZIP9 subfamily (SLC39A9) encodes a protein with characteristics of a membrane androgen receptor (mAR). Human ZIP9 is a 307-amino acid protein, which has eight putative transmembrane domains. ZIP9 has also been identified and characterized as a resident protein in the Golgi in DT40 and HeLa cell lines (PMID: 19420709, 23505453). ZIP9 may be present on the cell surface, as well as nuclear and mitochondrial membranes, in human breast and prostate cancer cell lines, where it mediates testosterone induction of apoptosis (PMID: 29952128).  $knock out of {\it ZIP9} \ would \ lead \ to \ zinc \ accumulation \ in \ the \ secretory \ pathway, \ which \ could \ influence \ the \ general \ ion$ homeostasis and in turn the glycosylation outcome (PMID: 36648436).

#### Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17607-1-AP (SLC39A9 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 17607-1-AP (SLC39A9 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).