

For Research Use Only

MYOM3 Polyclonal antibody

Catalog Number: 17692-1-AP

11 Publications



Basic Information

Catalog Number: 17692-1-AP	GenBank Accession Number: BC067101	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 µg/ml by Nanodrop and 267 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 127294	Recommended Dilutions: WB 1:500-1:3000 IP 0.5-4.0 ug for IP and 1:500-1:2000 for WB IHC 1:20-1:400
Source: Rabbit	Full Name: myomesin family, member 3	
Isotype: IgG	Calculated MW: 162 kDa, 136 kDa	
Immunogen Catalog Number: AG11849	Observed MW: 162 kDa	

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

ELISA, IF, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, canine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse skeletal muscle tissue, rat skeletal muscle tissue

IP : mouse skeletal muscle tissue,

IHC : human skeletal muscle tissue, mouse skeletal muscle tissue

Background Information

MYOM3 (myomesin 3) is a structural component of the M-band in striated muscle and is involved in sarcomere stability and resistance during intense or sustained stretching. MYOM3 can be detected mainly in intermediate speed fibers of skeletal muscle. Recently high level of MYOM3 fragments were detected in sera from patients with muscular dystrophy, including Duchenne muscular dystrophy (DMD). MYOM3 fragments may be used as serum biomarker for DMD and other neuromuscular disorders. This antibody recognizes the intact MYOM3 protein (160-160 kDa) as well as MYOM3 fragments (100 kDa and 130 kDa). (26060189)

Notable Publications

Author	Pubmed ID	Journal	Application
Minjung Lee	33124011	Int J Sports Med	ELISA
Flavien Bizot	36514350	Mol Ther Nucleic Acids	WB
Jeffrey M Lynch	30395621	PLoS One	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 17692-1-AP (MYOM3 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 17692-1-AP (MYOM3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




IP result of anti-MYOM3(IP:17692-1-AP, 4ug; Detection:17692-1-AP 1:1000) with mouse skeletal muscle tissue lysate 1280 ug.




Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue slide using 17692-1-AP (MYOM3 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue slide using 17692-1-AP (MYOM3 Antibody) at dilution of 1:200 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 17692-1-AP (MYOM3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 17692-1-AP (MYOM3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).