For Research Use Only

ABL2 Polyclonal antibody

Catalog Number: 17693-1-AP

Featured Product

12 Publications



Purification Method:

WB 1:1000-1:4000

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

Basic Information

Catalog Number: 17693-1-AP

Size:

BC065912

150ul, Concentration: 600 µg/ml by Nanodrop and 253 µg/ml by Bradford Full Name:

method using BSA as the standard;

Rabbit

Isotype: IgG Immunogen Catalog Number:

AG11860

GenBank Accession Number:

GeneID (NCBI):

v-abl Abelson murine leukemia viral protein lysate IHC 1:50-1:500 oncogene homolog 2 (arg, Abelson-

related gene) Calculated MW: 1167 aa, 127 kDa

Observed MW: 128-140 kDa

Applications

Tested Applications: IHC. IP. WB. FLISA **Cited Applications:**

CoIP, IF, IHC, WB Species Specificity: human, mouse, rat

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse brain tissue,

IP: HeLa cells.

IHC: human gliomas tissue,

Background Information

ABL2(Abelson tyrosine-protein kinase 2) is also named as ABLL, ARG and belongs to the protein kinase superfamily. It is a membrane-anchored nonreceptor tyrosine kinase that is a novel mediator of SEMA3F-induced RhoA inactivation and collapsing activity and it has a role in cytoskeletal rearrangements by its C-terminal F-actin- and microtubule-binding sequences. ABL2 can be polyubiquitinated and the polyubiquitination of ABL2 leads to degradation. This protein has 9 isoforms produced by alternative splicing with the molecular weight from 114 kDa to 128 kDa. The antibody is specific to the others except the 60 kDa isoform of ABL2.

Notable Publications

Author	Pubmed ID	Journal	Application
Xinkun Shen	34901556	Bioact Mater	IF
Siddhesh Aras	27913209	Biochim Biophys Acta	WB
Ze Zhang	34750518	Oncogene	WB,CoIP,IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 17693-1-AP (ABL2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 17693-1-AP (ABL2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP Result of anti-ABL2 (IP:17693-1-AP, 4ug; Detection:17693-1-AP 1:500) with HeLa cells lysate 1200ug.