

For Research Use Only

ALKBH2 Polyclonal antibody

Catalog Number: 17809-1-AP



Basic Information

Catalog Number: 17809-1-AP	GenBank Accession Number: BC070489	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 500 ug/ml by Nanodrop;	GeneID (NCBI): 121642	Recommended Dilutions: WB 1:500-1:1000
Source: Rabbit	UNIPROT ID: Q6NS38	
Isotype: IgG	Full Name: alkB, alkylation repair homolog 2 (E. coli)	
Immunogen Catalog Number: AG12108	Calculated MW: 261 aa, 29 kDa	
	Observed MW: 30-35 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : Caco-2 cells,
Species Specificity: human	

Background Information

Escherichia coli AlkB is a DNA repair enzyme that represents the founding member of the AlkB family of non-heme iron-dependent dioxygenases. The human genome encodes nine AlkB homologs termed AlkB homolog 1 through 8 (ALKBH1 through 8) plus a ninth member termed fat mass and obesity associated (FTO). Bacterial AlkB as well as human ALKBH2 and ALKBH3 catalyze the direct reversal of DNA methylation damage through a distinct oxidative demethylation reaction. Importantly, ALKBH2 has been shown to play a role in DNA alkylation damage repair in human, mouse and plant cells. Full-length ALKBH2 has a molecular weight of about 30kDa and ALKBH2 variant encoded by an alternatively spliced transcript with a frameshift deletion of the C-terminal dioxygenase domain has a molecular weight of about 25kDa (PMID:22079122).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17809-1-AP (ALKBH2 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.