#### For Research Use Only

# MGAT4C Polyclonal antibody

Catalog Number: 17841-1-AP

**Featured Product** 

2 Publications



**Purification Method:** 

WB 1:500-1:2000

protein lysate

Antigen affinity purification

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Recommended Dilutions:

**Basic Information** 

**Applications** 

Catalog Number:

17841-1-AP

Size:

150ul , Concentration: 300 µg/ml by Nanodrop and 187 µg/ml by Bradford UNIPROT ID:

method using BSA as the standard;

Source:

Rabbit Isotype:

Immunogen Catalog Number:

AG12201

**Tested Applications:** 

IP, WB, ELISA

**Cited Applications:** 

Species Specificity: human, mouse, rat Cited Species:

GenBank Accession Number:

BC064141

GeneID (NCBI): 25834

Q9UBM8 Full Name:

mannosyl (alpha-1,3-)-glycoprotein

beta-1.4-N-

acetylglucosaminyltransferase,

isozyme C (putative)

Calculated MW: 478 aa, 56 kDa

Observed MW: 55 kDa

Positive Controls:

WB: mouse testis tissue, rat testis tissue

IP: mouse testis tissue,

## **Background Information**

MGAT4C is predicted to be a glycosyltransferase that catalyzes a GlcNAc β1-4 linkage to the core mannose residues of N-glycans, resulting in N-glycans with tri- or tetraantennary structures. Recently MGAT4C has been found to get involved in processing complex N-glycan structures on CD133 and CD133 has been identified as a substrate of MGAT4C. Given to the fact that CD133 is widely used as a marker of numerous stem cell and cancer stem cell types, MGAT4C may have an important role in certain cancers. (PMID: 21937449)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Lijie Zhou	34185414	Clin Transl Med	WB
Mak Anthony B AB	21937449	J Biol Chem	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

### Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 17841-1-AP (MGAT4C antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-MGAT4C (IP:17841-1-AP, 4ug; Detection:17841-1-AP 1:400) with mouse testis tissue lysate 4000ug.