For Research Use Only

## GPR177 Polyclonal antibody Catalog Number: 17950-1-AP Featured Product 8 Pd

Featured Product 8 Publications





Basic Information	Catalog Number: 17950-1-AP	GenBank Accession Number BC110826	<ul> <li>Purification Method:</li> <li>Antigen affinity purification</li> </ul>
	Size: 150ul, Concentration: 650 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG12428	GenelD (NCBI):	Recommended Dilutions:
		79971	WB 1:1000-1:5000
		UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
		Q5T9L3	IHC 1:50-1:500
		Full Name: G protein-coupled receptor 177 Calculated MW:	
		Observed MW: 46-65 kDa	
		Applications	Tested Applications:
WB, IP, IHC, ELISA Cited Applications:			U-87 MG cells, mouse brain tissue, HEK-293 cell rain tissue
WB, IHC	HC IP : mouse liver tissue,		
Species Specificity: human, mouse, rat			: human ovary tissue, mouse liver tissue, humar ey tissue
Cited Species: human, goat, rat, mouse	KIGH		
TE buffer pH 9.0; (*) Alternati retrieval may be performed w buffer pH 6.0			
Background Information	Wnt protein secretion. GPR177 locali: apparatus, endosomes, and plasma n and gastric cancer. Several isoforms and 51 kDa, respectively. The sequer	zes to compartments of the se nembrane. GPR177 is overexp of GPR177 exist due to the alt ace can be processed by remo nigragrating band above 62 l	cretory pathway, including the ER, Golgi pressed in several cancer types including glioma ernative splicing, with predicted MW around 62
	Wnt protein secretion. GPR177 locali: apparatus, endosomes, and plasma n and gastric cancer. Several isoforms and 51 kDa, respectively. The sequer form of 45-46 kDa. In addition, slow n modified by N-linked glycosylation (	zes to compartments of the se nembrane. GPR177 is overexp of GPR177 exist due to the alt ace can be processed by remo nigragrating band above 62 l	cretory pathway, including the ER, Golgi pressed in several cancer types including glioma ernative splicing, with predicted MW around 62 ving the signal peptide, which generates a matu
	Wnt protein secretion. GPR177 locali: apparatus, endosomes, and plasma n and gastric cancer. Several isoforms and 51 kDa, respectively. The sequer form of 45-46 kDa. In addition, slow n modified by N-linked glycosylation (	zes to compartments of the se nembrane. GPR177 is overexp of GPR177 exist due to the alt ice can be processed by remo nigragrating band above 62 k PMID: 20549736).	cretory pathway, including the ER, Golgi pressed in several cancer types including glioma emative splicing, with predicted MW around 62 ving the signal peptide, which generates a matu Da can also be detected because GPR177 is
	Wnt protein secretion. GPR177 locali:         apparatus, endosomes, and plasma m         and gastric cancer. Several isoforms         and 51 kDa, respectively. The sequer         form of 45-46 kDa. In addition, slow m         modified by N-linked glycosylation (         Author       Pub         Qun Niu       345	tes to compartments of the se hembrane. GPR177 is overexp of GPR177 exist due to the alt ice can be processed by remo- nigragrating band above 62 k PMID: 20549736). med ID Journal	pressed in several cancer types including glioma emative splicing, with predicted MW around 62 ving the signal peptide, which generates a matu (Da can also be detected because GPR177 is Application IHC
	Wnt protein secretion. GPR177 locali: apparatus, endosomes, and plasma m and gastric cancer. Several isoforms and 51 kDa, respectively. The sequer form of 45-46 kDa. In addition, slow m modified by N-linked glycosylation (AuthorPub Qun NiuQun Niu345Shuyu Hu339	tes to compartments of the se hembrane. GPR177 is overexp of GPR177 exist due to the alt icce can be processed by remoin ingragrating band above 62 k PMID: 20549736). med ID Journal 42891 Reprod Sci	Application HC HC HC HC HC HC HC HC HC HC
	Wnt protein secretion. GPR177 locali: apparatus, endosomes, and plasma m and gastric cancer. Several isoforms and 51 kDa, respectively. The sequer form of 45-46 kDa. In addition, slow m modified by N-linked glycosylation (AuthorPub Qun NiuQun Niu345Shuyu Hu339	tes to compartments of the se hembrane. GPR177 is overexp of GPR177 exist due to the alt ice can be processed by remoin ingragrating band above 62 k PMID: 20549736). med ID Journal 42891 Reprod Sci 83548 Int Ophthalm	Application HCC HCC HCC HCC HCC HCC HCC HC
Notable Publications	Wnt protein secretion. GPR177 locali: apparatus, endosomes, and plasma m and gastric cancer. Several isoforms and 51 kDa, respectively. The sequer form of 45-46 kDa. In addition, slow m modified by N-linked glycosylation (AuthorPub Qun NiuQun Niu345Shuyu Hu339	res to compartments of the se hembrane. GPR177 is overexp of GPR177 exist due to the alt ice can be processed by remo- migragrating band above 62 k PMID: 20549736). med ID Journal 42891 Reprod Sci 83548 Int Ophthalm 69011 Front Physio	Application IHC Model MB MB MB MB MB MB MB MB MB MB MB MB MB M
Background Information Notable Publications Storage	Wnt protein secretion. GPR177 locali:         apparatus, endosomes, and plasma m         and gastric cancer. Several isoforms         and 51 kDa, respectively. The sequer         form of 45-46 kDa. In addition, slow m         modified by N-linked glycosylation (         Author       Pub         Qun Niu       345         Shuyu Hu       339         Yue Gao       335         Storage:       Storage Buffer:	res to compartments of the se hembrane. GPR177 is overexp of GPR177 exist due to the alt ice can be processed by remo- migragrating band above 62 k PMID: 20549736). med ID Journal 42891 Reprod Sci 83548 Int Ophthalm 69011 Front Physio er shipment. % glycerol pH 7.3.	Application HCC HCC HCC HCC HCC HCC HCC HC
Notable Publications	Wnt protein secretion. GPR177 locali:         apparatus, endosomes, and plasma m         and gastric cancer. Several isoforms         and 51 kDa, respectively. The sequer         form of 45-46 kDa. In addition, slow m         modified by N-linked glycosylation (         Author       Pub         Qun Niu       345         Shuyu Hu       339         Yue Gao       335         Storage:       Storage Buffer:         PBS with 0.02% sodium azide and 50	res to compartments of the se hembrane. GPR177 is overexp of GPR177 exist due to the alt ice can be processed by remo- migragrating band above 62 k PMID: 20549736). med ID Journal 42891 Reprod Sci 83548 Int Ophthalm 69011 Front Physio er shipment. % glycerol pH 7.3.	Application IHC Model MB MB MB MB MB MB MB MB MB MB MB MB MB M

and original va T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com


clusively available under Protein I his pro Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





WB result of GPR177 antibody (17950-1-AP; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GPR177 transfected HEK-293 cells. Various cell lysates were subjected to SDS PAGE followed by western blot with 17950-1-AP (GPR177 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



IP result of anti-GPR177 (IP:17950-1-AP, 3ug; Detection:17950-1-AP 1:500) with mouse liver tissue lysate 12500ug.



Immunohistochemical analysis of paraffinembedded human ovary using 17950-1-AP (GPR177 antibody) at dilution of 1:50 (under 10x lens).