## For Research Use Only

# RGS9 Polyclonal antibody

Catalog Number: 17970-1-AP

1 Publications



**Basic Information** 

**Applications** 

Catalog Number: 17970-1-AP

GenBank Accession Number: BC022504

**Purification Method:** Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

WB 1:500-1:3000

150ul, Concentration: 650 µg/ml by

8787 Nanodrop and 367 µg/ml by Bradford Full Name:

regulator of G-protein signaling 9

method using BSA as the standard;

Calculated MW: 622 aa, 71 kDa

Isotype: IgG

Observed MW:

58 kDa, 70-75 kDa

Immunogen Catalog Number:

AG11976

Rabbit

**Positive Controls:** 

**Tested Applications:** WB,ELISA

**Cited Applications:** 

Species Specificity: human, mouse, rat

**Cited Species:** 

human

WB: mouse brain tissue, SH-SY5Y cells

## **Background Information**

RGS9 is a member of the RGS family of signaling proteins that suppress the activity of G proteins by promoting their deactivation. There are two splice isoforms of RGS9 with quite different properties and patterns of expression. RGS9-1 is mainly found in the eye and is involved in regulation of phototransduction in rod and cone cells of the retina, while RGS9-2 is is expressed specifically in the striatum, a brain region involved in controlling movement, motivation, mood and addition. This antibody got about 58kD in the mouse eye and 70kD in the mouse brain in western blotting, both in line with their expected masses. Defects in RGS9 are a cause of prolonged electroretinal response suppression (PERRS).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Zhuo Shao	34573334	Genes (Basel)	WB

### Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 17970-1-AP (RGS9 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.