### For Research Use Only

# MLN51/CASC3 Polyclonal antibody

Catalog Number: 18047-1-AP

2 Publications



**Basic Information** 

Catalog Number: 18047-1-AP

GenBank Accession Number: BC050526

**Purification Method:** Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 650 µg/ml by

22794

WB 1:500-1:1000

Nanodrop and 333 µg/ml by Bradford Full Name:

cancer susceptibility candidate 3

IHC 1:50-1:500 IF 1:10-1:100

method using BSA as the standard;

Calculated MW: Rabbit 703 aa, 76 kDa

Isotype: Observed MW: IgG 115-120 kDa

Immunogen Catalog Number:

AG12694

**Applications** 

**Tested Applications:** 

IF, IHC, WB, ELISA

**Cited Applications:** 

IF, WB

Species Specificity: human

**Cited Species:** 

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: A2780 cells, MCF-7 cells, SKOV-3 cells

IHC: human liver cancer tissue,

IF: HepG2 cells,

## **Background Information**

CASC3(cancer susceptibility candidate 3) is also named as MLN51 and belongs to the CASC3 family. It is induced by CSF2, and is important in the proliferation of RA fibroblast-like synoviocytes in the pathogenesis of rheumatoid arthritis (RA)(PMID:17101062). It is a nucleocytoplasmic protein containing, within its aminoterminal half, a coiledcoil domain followed by two nuclear localization signals (NLSs) responsible for nuclear localization and, in its carboxyl-terminal half, a nuclear export signal (NES) that mediates cytoplasmic retention. It possesses many potential phosphorylation sites and is found to be phosphorylated in both normal and cancer-derived cells. (PMID:17652158).

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Ivone G Bruno	21596314	Mol Cell	WB
Daguenet Elisabeth E	22419818	Mol Biol Cell	WB,IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 18047-1-AP (MLN51/CASC3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 18047-1-AP (MLN51/CASC3 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells, using CASC3 antibody 18047-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).