### For Research Use Only

# TRAF3 Polyclonal antibody

Catalog Number: 18099-1-AP

**Featured Product** 

19 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

18099-1-AP BC075087 GeneID (NCBI):

150ul, Concentration: 800 µg/ml by 7187

Nanodrop;

Source: TNF receptor-associated factor 3

Rabbit Calculated MW: Isotype: 568 aa, 64 kDa IgG Observed MW: Immunogen Catalog Number: 55-60 kDa

AG12185

**Tested Applications:** IP, WB, ELISA

**Cited Applications:** 

IF, WB

Species Specificity: human, mouse

Cited Species:

human, rat, sheep, mouse, pig

**Purification Method:** 

Antigen affinity purification

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate

WB 1:1000-1:6000

**Applications** 

Positive Controls:

WB: COLO 320 cells, HEK-293 cells, multi-cells, HeLa

cells, Raji cells, NIH/3T3 cells

IP: Raji cells,

# **Background Information**

TNF receptor-associated factor 3 (TRAF3) has many alternative names including CAP-1, CD40 receptor-associated factor 1(CRAF1), CD40-binding protein(CD40BP) and LMP1-associated protein 1 (LAP1). TRAF3 regulates pathways leading to the activation of NF-kappa-B and MAP kinases, and plays a central role in the regulation of innate immune response. It modulates signaling pathways leading to the production of cytokines and interferon. As an essential constituent of several E3 ubiquitin-protein ligase complexes, TRAF3 promote 'Lys-63'-linked  $ubiquitination \ of \ target\ proteins.\ Moreover, TRAF3\ undergoes\ both\ 'Lys-48'-linked\ and\ 'Lys-63'-linked\ ubiquitination$ in response to various stimuli and is deubiquitinated by OTUB1, OTUB2 and OTUD5. Several alternatively spliced transcript variants encoding distinct isoforms (64 kDa and 55 kDa) have been reported.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Hai-Wei Chen	26456228	Sci Rep	WB
Yanli Wei	33028715	J Virol	WB
Jun Kang	33148796	J Virol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 18099-1-AP (TRAF3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



WB result of TRAF3 antibody (18099-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TRAF3 transfected HEK-293 cells.



IP result of anti-TRAF3 (IP:18099-1-AP, 4ug; Detection:18099-1-AP 1:1000) with Raji cells lysate 3160 ug.



Various lysates were subjected to SDS PAGE followed by western blot with 18099-1-AP (TRAF3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.