

For Research Use Only

FARSA Polyclonal antibody

Catalog Number: 18121-1-AP

Featured Product

5 Publications



Basic Information

Catalog Number: 18121-1-AP	GenBank Accession Number: BC006495	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 1000 µg/ml by Nanodrop and 400 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2193	Recommended Dilutions: WB 1:4000-1:16000 IP 0.5-4.0 ug for IP and 1:1000-1:4000 for WB
Source: Rabbit	Full Name: phenylalanyl-tRNA synthetase, alpha subunit	IHC 1:500-1:2000 IF 1:10-1:100
Isotype: IgG	Calculated MW: 58 kDa	
Immunogen Catalog Number: AG12647	Observed MW: 57 kDa	

Applications

Tested Applications:
IF, IHC, IP, WB, ELISA

Cited Applications:
ELISA, IF, IP, RIP, WB

Species Specificity:
human, mouse

Cited Species:
human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, K-562 cells, MCF-7 cells, HepG2 cells

IP : MCF-7 cells,

IHC : human liver cancer tissue, human kidney tissue

IF : MCF-7 cells, Hela cells

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Yoshinao Muro	36191779	Autoimmun Rev	WB,IP
Megha Jhanji	35688816	Nat Commun	WB
Qian Zhou	35879296	Nat Commun	IF,ELISA

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18121-1-AP (FARSA antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




IP result of anti-FARS (IP:18121-1-AP, 4ug; Detection:18121-1-AP 1:2000) with MCF-7 cells lysate 3240 ug.



Immunofluorescent analysis of MCF-7 cells, using FARSA antibody 18121-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 18121-1-AP (FARSA antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 18121-1-AP (FARSA antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).