For Research Use Only

UBE4B Polyclonal antibody Catalog Number:18148-1-AP Featured Product

Featured Product 7 Publications





Basic Information	Catalog Number: 18148-1-AP			Purification Method: Antigen affinity purification			
	Size:	GeneID (NCBI): Recoil 10277 WB 1 UNIPROT ID: IP 0.5 095155 prote Full Name: IHC 1		Recommended Dilutions: WB 1:2000-1:10000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:20-1:200 IF/ICC 1:10-1:100			
	150ul , Concentration: 400 ug/ml by Nanodrop and 247 ug/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG12836						
						Observed MW:	
146 kDa							
Applications				Tested Applications:	Positive Controls:		
		WB, IHC, IF/ICC, IP, ELISA		WB : HEK-293T cells, mouse testis tissue, HeLa cells, K 562 cells, Raji cells IP : HeLa cells,			
		Cited Applications: WB, IHC					
	Species Specificity:						
	human, mouse, rat			an heart tissue,			
	Cited Species: human	IF/ICC : HeLa cells,					
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0						
Background Information	UBE4B (Ubiquitin Conjugation E4 B) is a protein-coding gene that plays a crucial role in the ubiquitination process, which is essential for protein degradation and regulation of various cellular processes. UBE4B functions as both an E3 and E4 ubiquitin ligase. It collaborates with other ubiquitin ligases to catalyze the transfer of ubiquitin molecules to target proteins, leading to their degradation or other intracellular sorting events. For example, UBE4B has been shown to ubiquitinate and degrade proteins such as Ku70, c-FLIPL, and p53. It also interacts with the E3 ubiquitin ligase ITCH to induce polyubiquitination and proteasomal degradation of target proteins.						
	molecules to target proteins, leading has been shown to ubiquitinate and d	legrade proteins such as l	Ku70, c-FLIPL,	ular sorting events. For example, UBE4B and p53. It also interacts with the E3			
Notable Publications	molecules to target proteins, leading has been shown to ubiquitinate and d ubiquitin ligase ITCH to induce polyu	legrade proteins such as l	Ku70, c-FLIPL, somal degrad	ular sorting events. For example, UBE4B and p53. It also interacts with the E3 dation of target proteins.			
Notable Publications	molecules to target proteins, leading has been shown to ubiquitinate and d ubiquitin ligase ITCH to induce polyu	legrade proteins such as l biquitination and protea	〈u70, c-FLIPL, somal degrad	ular sorting events. For example, UBE4B and p53. It also interacts with the E3			
Notable Publications	molecules to target proteins, leading has been shown to ubiquitinate and d ubiquitin ligase ITCH to induce polyu Author Pub Xianyan Hu 326	legrade proteins such as I biquitination and protea med ID Journal	Ku70, c-FLIPL, somal degrad	ular sorting events. For example, UBE4B and p53. It also interacts with the E3 dation of target proteins. Application			
Notable Publications	molecules to target proteins, leading has been shown to ubiquitinate and d ubiquitin ligase ITCH to induce polyu Author Pub Xianyan Hu 326 Hongyu Zhang 249	legrade proteins such as I biquitination and protea med ID Journal 45369 Mol Cell 95952 Proteom	Ku70, c-FLIPL, somal degrad	ular sorting events. For example, UBE4B and p53. It also interacts with the E3 dation of target proteins. Application WB			
	molecules to target proteins, leading has been shown to ubiquitinate and d ubiquitin ligase ITCH to induce polyu Author Pub Xianyan Hu 326 Hongyu Zhang 249 Xu-Qiong Huang 320 Storage: Stora at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°	egrade proteins such as I biquitination and protea med ID Journal 45369 Mol Cell 95952 Proteom 21266 Onco Ta er shipment. % glycerol pH 7.3.	Ku70, c-FLIPL, somal degrad l	ular sorting events. For example, UBE4B and p53. It also interacts with the E3 dation of target proteins. Application WB WB			
Notable Publications Storage	molecules to target proteins, leading has been shown to ubiquitinate and d ubiquitin ligase ITCH to induce polyu Author Pub Xianyan Hu 326 Hongyu Zhang 249 Xu-Qiong Huang 320 Storage: Storage: Storage Buffer:	egrade proteins such as I biquitination and protea med ID Journal 45369 Mol Cell 95952 Proteom 21266 Onco Ta er shipment. % glycerol pH 7.3.	Ku70, c-FLIPL, somal degrad l	ular sorting events. For example, UBE4B and p53. It also interacts with the E3 dation of target proteins. Application WB WB			

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data









Immunohistochemical analysis of paraffinembedded human heart tissue slide using 18148-1-AP (UBE4B Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 18148-1-AP (UBE4B Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HeLa cells using 18148-1-AP (UBE4B antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.



IP result of anti-UBE4B (IP:18148-1-AP, 4ug; Detection:18148-1-AP 1:1000) with HeLa cells lysate 2320 ug.