For Research Use Only

## PD-L2/CD273 Polyclonal antibody

Catalog Number:18251-1-AP 17 Publications




## Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 18251-1-AP BC074766 Antigen affinity purification GenelD (NCBI): Recommended Dilutions: Size: 150ul , Concentration: 400 ug/ml by 80380 WB 1:500-1:2000 Nanodrop: UNIPROT ID: Source: Q9BQ51 Rabbit Full Name: Isotype: programmed cell death 1 ligand 2 lgG Calculated MW: Immunogen Catalog Number: 273 aa, 31 kDa AG13039 **Observed MW:** 40-50 kDa, 28 kDa **Applications Tested Applications: Positive Controls:** WB, ELISA WB: NIH/3T3 cells, mouse heart tissue, HepG2 cells, **Cited Applications:** RAW 264.7 cells WB, IHC **Species Specificity:** human, mouse, rat **Cited Species:** human, mouse **Background Information** Programmed cell death ligand 2 (PD-L2, also known as CD273, or B7-DC), is one of the B7 family molecules which are type I transmembrane proteins belonging to the immunoglobulin superfamily. PD-L2 was identified both by its homology to PD-L1 and by analysis of genes differentially expressed in libraries generated from dendritic cells (DC) and activated macrophages (PMID: 14515254). PD-L2 is a second ligand for PD-1, a costimulatory molecule that plays an inhibitory role in regulating T cell activation in the periphery. **Notable Publications** Author Pubmed ID Journal Application IHC Xiuliu 32969157 1 Cell Mol Med WB,IHC Qixia Xu 31493351 Aging Cell Xianlong Chen 36325339 Front Immunol IHC Storage Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol, pH7.3 Aliquoting is unnecessary for -20°C storage \*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 18251-1-AP (PD-L2/CD273 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.