

For Research Use Only

# STRAP Polyclonal antibody

Catalog Number: 18277-1-AP

Featured Product

9 Publications



## Basic Information

### Catalog Number:

18277-1-AP

### Size:

150ul, Concentration: 450 ug/ml by Nanodrop and 307 ug/ml by Bradford method using BSA as the standard;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG13085

### GenBank Accession Number:

BC000162

### GeneID (NCBI):

11171

### UNIPROT ID:

Q9Y3F4

### Full Name:

serine/threonine kinase receptor associated protein

### Calculated MW:

38 kDa

### Observed MW:

39 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:1000-1:6000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

### Cited Applications:

WB, IHC, IF, IP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, hamster

### Positive Controls:

WB : HeLa cells, HEK-293 cells, mouse brain tissue, rat brain tissue

IP : HeLa cells,

IHC : human ovary tumor tissue, mouse testis tissue

IF/ICC : HeLa cells,

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Background Information

STRAP (Serine-threonine kinase receptor-associated protein), also known as MAWD, is a ubiquitous WD40 domain protein, common function of which is to provide a suitable scaffold for coordinating multiprotein complex assemblies and thus, regulate a variety of cellular processes like signal transduction, transcriptional regulation, programmed cell death and so on. STRAP plays a role in the cellular distribution of the SMN complex, which is essential for spliceosomal snRNP assembly in the cytoplasm. STRAP inhibits transforming growth factor-beta (TGF-beta) signaling and enhances tumorigenicity via TGF-beta-dependent and -independent mechanisms. STRAP is up-regulated in several cancers and functions as an oncogene. STRAP imparts oncogenic characteristics to cells by promoting ERK and pRb phosphorylation. Moreover, STRAP regulates c-Jun stability by decreasing the ubiquitylation and proteosomal degradation of c-Jun.

## Notable Publications

Author	Pubmed ID	Journal	Application
Shanshan Hu	33134183	Front Oncol	WB
Evan P Booy	30247708	Nucleic Acids Res	WB,IP
Muhammad Riaz Khan	29046333	EMBO J	WB,IP

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18277-1-AP (STRAP antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




WB result of STRAP antibody (18277-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-STRAP transfected HEK-293 cells.



Immunohistochemical analysis of paraffin-embedded human ovary tumor tissue slide using 18277-1-AP (STRAP antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-STRAP (IP:18277-1-AP, 4ug; Detection:18277-1-AP 1:2000) with HeLa cells lysate 2000ug.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using STRAP antibody (18277-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L).