#### For Research Use Only

# G-CSFR Polyclonal antibody

Catalog Number: 18310-1-AP

2 Publications



**Basic Information** 

Catalog Number: 18310-1-AP

method using BSA as the standard;

Size:

Rabbit

Isotype:

GenBank Accession Number: BC053585

DC023303

GeneID (NCBI):

150ul , Concentration: 300 µg/ml by 1441

Nanodrop and 200 µg/ml by Bradford Full Name:

colony stimulating factor 3 receptor

(granulocyte)

Calculated MW:

95 kDa Observed MW:

85 kDa

**Applications** 

Tested Applications:

FC, WB,ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

human, mouse

Recommended Dilutions:

**Purification Method:** 

WB 1:500-1:2000

Antigen affinity purification

Positive Controls:

WB: human placenta tissue,

### **Background Information**

G-CSF-R, also named as CSF3R, CD114 and GCSFR is the receptor for colony stimulating factor 3, a cytokine that controls the production, differentiation, and function of granulocytes. It is a 130 kD to 150 kD glycoprotein single chain receptor which binds and activated by GCSF. In addition it may function in some adhesion or recognition events at the cell surface. It has been classified as a member of the hematopoietic (cytokine) receptor family, cytokine receptor class I, or the gp 130 related cytokine receptor family (although it does not apparently bind to gp 130). GCSF receptors can be found on neutrophils, myeloid leukemia cells that respond to GCSF, bone marrow cells of neutrophilic granulocyte lineage, and on placental trophoblasts. In addition, a soluble form is also expressed. Mutations in this gene are a cause of Kostmann syndrome, also known as severe congenital neutropenia.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Mary P Udumula	34144215	Mol Metab	WB
Yang Cheng	30779348	J Cell Physiol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



human placenta tissue were subjected to SDS PAGE followed by western blot with 18310-1-AP (G-CSFR antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



1X10^6 U-937 cells were stained with 0.2ug G-CSFR antibody (18310-1-AP, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.