For Research Use Only

MAD1 Polyclonal antibody

Catalog Number: 18322-1-AP

5 Publications



Basic Information

Catalog Number: GenBank Accession Number: 18322-1-AP

BC009964

GeneID (NCBI):

150ul, Concentration: 800 µg/ml by 8379

Nanodrop; Full Name:

Source: MAD1 mitotic arrest deficient-like 1

Rabbit

Isotype: Calculated MW:

83 kDa IgG Observed MW:

Immunogen Catalog Number:

83 kDa AG13171

Applications

Tested Applications:

IF, IHC, IP, WB, ELISA

Cited Applications:

IF, WB

Species Specificity:

human **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method:

Antigen affinity purification

Recommended Dilutions: WB 1:2000-1:16000

IP 0.5-4.0 ug for IP and 1:500-1:2000

for WB IHC 1:20-1:200 IF 1:50-1:500

Positive Controls:

WB: A431 cells, human brain tissue, Raji cells, HeLa

cells, HepG2 cells

IP: HeLa cells.

IHC: human brain tissue, human kidney tissue, human ovary tissue, human placenta tissue, human skin tissue, human spleen tissue, human testis tissue

IF: A431 cells, SH-SY5Y cells

Background Information

MAD1L1 is a component of the spindle-assembly checkpoint that prevents the onset of anaphase until all chromosomes are properly aligned at the metaphase plate. It recruits MAD2L1 to unattached kinetochores. Also it is required for anchoring MAD2L1 to the nuclear periphery. In addition, it binds to the TERT promoter and represses telomerase expression, possibly by interfering with MYC binding [PMID:10049595, 12837246].

Notable Publications

Author	Pubmed ID	Journal	Application
Carolina Villarroya-Beltri	36322655	Sci Adv	WB
Fengjie Yuan	31291454	Nucleic Acids Res	WB,IF
Sara M Reed	24621507	Cell Cycle	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18322-1-AP (MAD1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



IP Result of anti-MAD1L1 (IP:18322-1-AP, 5ug; Detection:18322-1-AP 1:1000) with HeLa cells lysate 880ug.



Immunohistochemical analysis of paraffinembedded human brain using 18322-1-AP (MAD1 antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using MAD1 antibody (18322-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).