For Research Use Only

## CD109 Polyclonal antibody

Catalog Number:18715-1-AP

1 Publications





Basic Information	Catalog Number: 18715-1-AP	GenBank Accession Number: BC 152996	Purification Method: Antigen affinity purification
	Size: 150ul , Concentration: 273 µg/ml by	GenelD (NCBI): 135228	Recommended Dilutions: WB 1:500-1:1000
	Bradford method using BSA as the standard;	UNIPROT ID: O6YHK3	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
	Source: Rabbit	Full Name: CD109 molecule	
	lsotype: IgG	Calculated MW: 162 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IP, ELISA	WB: PC-3 cells,	
	Cited Applications: WB	IP : PC-3 cells,	
	Species Specificity: human		
	Cited Species: human		
Background Information	cell surface of platelets, activated T- of transforming growth factor beta (T human keratinocytes. Up regulation tumor.CD109 is a glycosyl-phosphati	inked cell surface glycoprotein of a cells, and endothelial cells. CD109 GF-beta). CD109 has been identifi of CD109 expression has been obs dylinositol-anchored glycoprotein hioester-containing proteins. Ther	approximately 170kd which is found on the blinds to and negatively regulates signali ed as part of the TGF-beta receptor system erved in several different types of a that is a member of the a2- re're 2 forms of 150 (p150) and 120 kDa (p1
	cell surface of platelets, activated T- of transforming growth factor beta (T human keratinocytes. Up regulation t tumor.CD109 is a glycosyl-phosphati macroglobulin/C3, C4, C5 family of t exist due to proteolytic degradation	inked cell surface glycoprotein of a cells, and endothelial cells. CD109 GF-beta). CD109 has been identifi of CD109 expression has been obs dylinositol-anchored glycoprotein hioester-containing proteins. Ther	approximately 170kd which is found on the 9 binds to and negatively regulates signali ed as part of the TGF-beta receptor system erved in several different types of 1 that is a member of the a2- re're 2 forms of 150 (p150) and 120 kDa (p1
Background Information	cell surface of platelets, activated T- of transforming growth factor beta (T human keratinocytes. Up regulation of tumor.CD109 is a glycosyl-phosphati macroglobulin/C3, C4, C5 family of t exist due to proteolytic degradation f Author Pul	inked cell surface glycoprotein of a cells, and endothelial cells. CD109 GF-beta). CD109 has been identifi of CD109 expression has been obs dylinositol-anchored glycoprotein hioester-containing proteins. Ther from a 180 kDa form. The antibody	approximately 170kd which is found on the binds to and negatively regulates signali ed as part of the TGF-beta receptor system erved in several different types of 1 that is a member of the a2- re're 2 forms of 150 (p150) and 120 kDa (p1 r can recognize all these 4 isoforms.
	cell surface of plateletts, activated T- of transforming growth factor beta (T human keratinocytes. Up regulation of tumor.CD109 is a glycosyl-phosphati macroglobulin/C3, C4, C5 family of t exist due to proteolytic degradation f Author Pul	inked cell surface glycoprotein of a cells, and endothelial cells. CD109 GF-beta). CD109 has been identifio of CD109 expression has been obs dylinositol-anchored glycoprotein hioester-containing proteins. Ther from a 180 kDa form. The antibody <b>bmed ID Journal</b> 113239 Oncol Lett	approximately 170kd which is found on the blinds to and negatively regulates signali ed as part of the TGF-beta receptor system erved in several different types of a that is a member of the a2- re're 2 forms of 150 (p150) and 120 kDa (p1 or can recognize all these 4 isoforms.

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





PC-3 cells were subjected to SDS PAGE followed by western blot with 18715-1-AP (CD109 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.

IP result of anti-CD109 (IP:18715-1-AP, 4ug; Detection:18715-1-AP 1:600) with PC-3 cells lysate 5000ug.