For Research Use Only

GPX5 Polyclonal antibody

Catalog Number: 18731-1-AP

8 Publications



Basic Information

Catalog Number: 18731-1-AP

GenBank Accession Number:

Purification Method:

NM 001509 GeneID (NCBI): Antigen affinity purification

Size 150ul, Concentration: 800 µg/ml by

Recommended Dilutions: WB 1:200-1:1000

Nanodrop and 267 µg/ml by Bradford Full Name:

IHC 1:20-1:200

method using BSA as the standard;

glutathione peroxidase 5 (epididymal

androgen-related protein)

Rabbit

Calculated MW:

Isotype: IgG

25 kDa Observed MW:

22-28 kDa

Applications

Tested Applications:

IHC, WB,ELISA

Cited Applications:

IF, WB

Positive Controls:

WB: mouse testis tissue, PC-3 cells, human testis

tissue, mouse liver tissue IHC: human testis tissue.

Species Specificity:

human, mouse, rat

Cited Species:

human, goat, mouse, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

GPX5 belongs to the glutathione peroxidase family. GPX5 protects cells and enzymes from oxidative damage, by catalyzing the reduction of hydrogen peroxide, lipid peroxides and organic hydroperoxide, by glutathione. Catalytic the reaction: 2 glutathione + H2O2 = glutathione disulfide + 2 H2O. GPX5 constitute a glutathionine peroxidase-like protective system against peroxide damage in sperm membrane lipids. GPX5 has a calculated molecular mass of 25 kDa and apparent molecular mass of 22-28 kDa due to the cleavage of signal peptide and glycosylation.

Notable Publications

Author	Pubmed ID	Journal	Application
Lin Huang	27003159	Sci Rep	IF
Ilias Michos	34204554	Animals (Basel)	WB
I Vilagran	25678437	Andrology	WB

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



mouse testis tissue were subjected to SDS PAGE followed by western blot with 18731-1-AP (GPXS antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis using 18731-1-AP (GPX5 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis using 18731-1-AP (GPX5 antibody) at dilution of 1:50 (under 40x lens).