#### For Research Use Only

# HSF4 Polyclonal antibody

Catalog Number: 18797-1-AP

Featured Product

4 Publications



**Basic Information** 

Catalog Number: 18797-1-AP

Nanodrop:

Source:

Rabbit

GenBank Accession Number:

BC130383

GeneID (NCBI):

Size: 150ul , Concentration: 500 ug/ml by

**UNIPROT ID:** Q9ULV5

Isotype: heat shock transcription factor 4

IgG Calculated MW:

53 kDa Observed MW:

Full Name:

53-58 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

Cited Applications:

WB, IHC

Species Specificity: human, mouse **Cited Species:** human, mouse

Positive Controls:

WB: HeLa cells, mouse skeletal muscle tissue, human skeletal muscle tissue, human heart tissue, MCF-7

**Purification Method:** 

WB 1:1000-1:4000

Antigen affinity purification

Recommended Dilutions:

### **Background Information**

HSF4 belongs to the HSF family. HSF4 is a DNA-binding protein that specifically binds heat shock promoter elements (HSE). The isoform HSF4A represses transcription while the isoform HSF4B activates transcription. HSF4 is a homotrimer which exhibits constitutive DNA binding and forms trimers even in the absence of stress. It is  $expressed in heart, skeletal \ muscle, eye\ and \ brain, and \ at \ much \ lower \ levels \ in \ some \ other \ tissues.$  Mutation of HSF4 will cause lamellar cataract which is known as Marner type cataract (CAM). Lamellar cataract is an autosomal dominant common type of infantile cataract. Finger malformation is observed in some kindreds. Inheritance of these forms of cataract is autosomal dominant.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Wenjing Zhang	36229759	Mol Cell Biochem	WB
Feng Guo	34975151	Oncogene	WB
Saeki Saito	40004241	Int J Mol Sci	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 18797-1-AP (HSF4 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.