For Research Use Only

BBS7 Polyclonal antibody

Catalog Number: 18961-1-AP

7 Publications



Basic Information

Catalog Number: 18961-1-AP

BC032691 GeneID (NCBI):

GenBank Accession Number:

Size: Genel D (150ul, Concentration: 900 µg/ml by 55212

Nanodrop and 453 µg/ml by Bradford Full Name:

method using BSA as the standard; Bardet-Biedl syndrome 7

Source: Calculated MW: 715 aa, 80 kDa Isotype: Observed MW: IgG 75 kDa

Immunogen Catalog Number:

AG4601

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:3000 IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB IHC 1:20-1:200

Applications

Tested Applications:

IHC, IP, WB, ELISA
Cited Applications:
IHC, IP, WB

Species Specificity: human, mouse Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: human brain tissue,
IP: mouse testis tissue,

IHC: human testis tissue, human brain tissue

Background Information

BBS7, also named as BBS2L1, is a component of the BBSome complex which is required for ciliogenesis but is dispensable for centriolar satellite function. BBS7 is mediated in part by the Rab8 GDP/GTP exchange factor, which localizes to the basal body and contacts the BBSome.

Notable Publications

Author	Pubmed ID	Journal	Application
Ying Hsu	29049287	PLoS Genet	WB
Poppy Datta	31694913	J Biol Chem	WB,IP
Satyabrata Sinha	24854858	Invest Ophthalmol Vis Sci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




human brain tissue were subjected to SDS PAGE followed by western blot with 18961-1-AP (BBS7 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis using 18961-1-AP (BBS7 antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis using 18961-1-AP (BBS7 antibody) at dilution of 1:50 (under 40x lens).



IP Result of anti-BBS7 (IP:18961-1-AP, 4ug; Detection:18961-1-AP 1:500) with mouse testis tissue lysate 4000ug.