

For Research Use Only

iNOS Polyclonal antibody

Catalog Number: 18985-1-AP

Featured Product

805 Publications



Basic Information

Catalog Number:

18985-1-AP

Size:

150ul, Concentration: 1200 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_000625

GeneID (NCBI):

4843

UNIPROT ID:

P35228

Full Name:

nitric oxide synthase 2, inducible

Calculated MW:

131 kDa

Observed MW:

110-130 kDa, 65-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:4000

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB, IF, FC (Intra), IP, ELISA

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, pig, chicken, bovine

Positive Controls:

WB : mouse liver tissue, rat liver tissue

IF/ICC : HepG2 cells, HeLa cells

Background Information

NOS2, also named as iNOS and NOS2A, produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. NO is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission, antimicrobial and antitumoral activities. NOS2 is a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. iNOS has a very short half-life due to rapid degradation by calpain. iNOS monomer is a direct substrate of calpain I and can be cleaved by calpain I at the canonical CaM-binding site(503-532aa) of iNOS, and then a ~70 kDa band can be detected by western (PMID:11786228). This antibody is specific to NOS2.

Notable Publications

Author	Pubmed ID	Journal	Application
Lin-Tao Xu	34601084	J Ethnopharmacol	WB
Yan Yang	34582523	Biomater Sci	IF,FC
Haoyu Guo	36201949	Biomaterials	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1%BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18985-1-AP (iNOS antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



WB result of iNOS antibody (18985-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-iNOS transfected HeLa cells.



1X10⁶ HepG2 cells were intracellularly stained with 0.4 ug Anti-Human iNOS (18985-1-AP) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using iNOS antibody (18985-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).