For Research Use Only

KIF18A Polyclonal antibody

Catalog Number:19245-1-AP

Featured Product 14 Publications





Basic Information	Catalog Number: 19245-1-AP	GenBank Accession Number: BC048347	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 700 ug/ml by		WB 1:500-1:2000	
	Nanodrop;	UNIPROT ID:	IHC 1:200-1:800	
	Source:	Q8NI77		
	Rabbit	Full Name:		
	Isotype: IgG	kinesin family member 18A		
	Immunogen Catalog Number: AG6166	Calculated MW: 102 kDa		
		Observed MW:		
		102 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, IHC, ELISA	WB : Jurkat cells, HeLa cells IHC : human colon cancer, human breast cancer tissue, human liver tissue		
	Cited Applications: WB, IHC, IF, IP			
	Species Specificity: human			
	Cited Species: human, mouse			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	hydrolysis. KIF18A is related to the limitenance of microtubule dynami	ocalization of membrane organelle cs and mitosis. KIF18A expression atocellular carcinoma and lung ade	d moves along them using energy from ATP as and protein complexes, as well as the is low in normal tissues and high in solid nocarcinoma, colorectal cancer, renal cell r weight of KIF18A is 102 kDa (PMID:	
	hydrolysis. KIF 18A is related to the l maintenance of microtubule dynami tumors, including breast cancer, hepa carcinoma, and head and neck squan 36830695).	ocalization of membrane organelle cs and mitosis. KIF18A expression atocellular carcinoma and lung ade	es and protein complexes, as well as the is low in normal tissues and high in solid nocarcinoma, colorectal cancer, renal cell r weight of KIF 18A is 102 kDa (PMID:	
	hydrolysis. KIF 18A is related to the li maintenance of microtubule dynami tumors, including breast cancer, hepa carcinoma, and head and neck squan 36830695).	ocalization of membrane organelle cs and mitosis. KIF18A expression atocellular carcinoma and lung ade nous cell carcinoma. The molecula	es and protein complexes, as well as the is low in normal tissues and high in solid nocarcinoma, colorectal cancer, renal cell	
Background Information	hydrolysis. KIF 18A is related to the li maintenance of microtubule dynami tumors, including breast cancer, hepa carcinoma, and head and neck squan 36830695). Author Put Fu-Tao Chen 317	ocalization of membrane organelle cs and mitosis. KIF 18A expression atocellular carcinoma and lung ade nous cell carcinoma. The molecula omed ID Journal 772692 Dis Markers	es and protein complexes, as well as the is low in normal tissues and high in solid nocarcinoma, colorectal cancer, renal cell r weight of KIF 18A is 102 kDa (PMID: Application WB	
	hydrolysis. KIF 18A is related to the L maintenance of microtubule dynami tumors, including breast cancer, hepa carcinoma, and head and neck squar 36830695). Author Put Fu-Tao Chen 317 Weijia Liao 254	ocalization of membrane organelle cs and mitosis. KIF18A expression atocellular carcinoma and lung ade nous cell carcinoma. The molecula	es and protein complexes, as well as the is low in normal tissues and high in solid nocarcinoma, colorectal cancer, renal cell r weight of KIF 18A is 102 kDa (PMID: Application	
Notable Publications	hydrolysis. KIF 18A is related to the L maintenance of microtubule dynami tumors, including breast cancer, hepa carcinoma, and head and neck squan 36830695). Author Put Fu-Tao Chen 317 Weijia Liao 254	ocalization of membrane organelle cs and mitosis. KIF 18A expression atocellular carcinoma and lung ade nous cell carcinoma. The molecula omed ID Journal 772692 Dis Markers i31949 Oncotarget i59823 Cell Div	es and protein complexes, as well as the is low in normal tissues and high in solid nocarcinoma, colorectal cancer, renal cell r weight of KIF 18A is 102 kDa (PMID: Application WB IHC	
	hydrolysis. KIF 18A is related to the U maintenance of microtubule dynami tumors, including breast cancer, hepa carcinoma, and head and neck squar 36830695). Author Put Fu-Tao Chen 317 Weijia Liao 254 Feng Tang 304 Storage:	ocalization of membrane organelle cs and mitosis. KIF 18A expression atocellular carcinoma and lung ade nous cell carcinoma. The molecula omed ID Journal 772692 Dis Markers 31949 Oncotarget 459823 Cell Div ter shipment. 9% glycerol pH 7.3.	es and protein complexes, as well as the is low in normal tissues and high in solid nocarcinoma, colorectal cancer, renal cell r weight of KIF 18A is 102 kDa (PMID: Application WB IHC	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Jurkat cells were subjected to SDS PAGE followed by western blot with 19245-1-AP (KIF18A antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human colon cancer slide using 19245-1-AP (KIF 18A antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).