For Research Use Only

CD57 Polyclonal antibody

Catalog Number: 19401-1-AP

5 Publications



Basic Information

Catalog Number: GenBank Accession Number: 19401-1-AP BC010466

Size: GeneID (NCBI):

150ul, Concentration: 300 µg/ml by Nanodrop and 267 µg/ml by Bradford

method using BSA as the standard;

Rabbit Calculated MW: 38 kDa Isotype: IgG Observed MW: Immunogen Catalog Number: 45 kDa, 70 kDa

AG13550

Applications

Tested Applications: IHC, WB, ELISA

Cited Applications:

IF, IHC

Species Specificity: human, mouse **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000 IHC 1:50-1:500

Positive Controls:

WB: PC-12 cells, SH-SY5Y cells

IHC: mouse brain tissue, mouse spleen tissue, human thyroid cancer tissue, human prostate cancer tissue

Background Information

B3GAT1 is associated with glycosaminoglycan biosynthesis. It plays a key role in neurodevelopment. The calculated molecular weight of B3GAT1 is 38 kDa and 19401-1-AP antibody detectes 45 and 70 kDa forms in SDS-PAGE which might as a result of different degrees of glycosylation or the formation of homodimer.

beta-1,3-glucuronyltransferase 1 (glucuronosyltransferase P)

Notable Publications

Author	Pubmed ID	Journal	Application
Sihui Yu	35626004	Cancers (Basel)	IHC
Jing Zhang	35042621	Gynecol Oncol	IHC
Jie Chen	34976977	Front Bioeng Biotechnol	IF

Storage


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 19401-1-AP (CD57 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 19401-1-AP (CD57 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).