# For Research Use Only

# ATG14/Barkor (N-terminal) Polyclonal antibody

www.ptglab.com

Catalog Number: 19491-1-AP

Featured Product

34 Publications

### **Basic Information**

Catalog Number: 19491-1-AP

Size:

GenBank Accession Number:

NM\_014924 GeneID (NCBI): 150ul , Concentration: 350 ug/ml by 22863

Nanodrop; **UNIPROT ID:** Source: Q6ZNE5 Rabbit

Full Name: Isotype: KIAA0831 IgG Calculated MW:

55 kDa Observed MW: 65-68 kDa

**Purification Method:** 

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:1000 IHC 1:50-1:500 IF/ICC 1:50-1:500

## **Applications**

**Tested Applications:** 

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IF, IP

Species Specificity: human, mouse

**Cited Species:** 

human, mouse, rat, pig, canine, bovine

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls:**

WB: HEK-293 cells, A431 cells, mouse brain tissue,

NIH/3T3 cells

IHC: mouse brain tissue, IF/ICC: MCF-7 cells,

# **Background Information**

Barkor, also named as KIAA0831, Atg14 and Atg14L, is required for both basal and inducible autophagy. It plays a role in autophagosome formation and MAP1LC3/LC3 conjugation to phosphatidylethanolamine. Barkor promotes BECN1 translocation from the trans-Golgi network to autophagosomes. This antibody is specific to KIAA0831(Barkor).

### **Notable Publications**

| Author               | Pubmed ID | Journal                           | Application |
|----------------------|-----------|-----------------------------------|-------------|
| Luis Muniz-Feliciano | 28933590  | Autophagy                         | WB          |
| Qiang Fu             | 25234643  | J Gen Virol                       | WB          |
| Xueqiang Peng        | 32949647  | Biochim Biophys Acta Mol Cell Res | WB          |

### Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


### **Selected Validation Data**



HEK-293 cells were subjected to SDS PAGE followed by western blot with 19491-1-AP (ATG14/Barkor (N-terminal) antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 19491-1-AP (ATG14/Barkor (N-terminal) antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using 19491-1-AP (ATG14/Barkor (N-terminal) antibody) at dilution of 1:50 and CoraLite488-Conjugated Goat Anti-Rabbit IgG(H+L).