

For Research Use Only

Aldehyde oxidase Polyclonal antibody

Catalog Number: 19495-1-AP

Featured Product

10 Publications

Basic Information

Catalog Number:

19495-1-AP

Size:

150ul, Concentration: 800 µg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_001159

GeneID (NCBI):

316

Full Name:

aldehyde oxidase 1

Calculated MW:

148 kDa

Observed MW:

148 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:200-1:1000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:50-1:500

IF 1:10-1:100

Applications

Tested Applications:

IF, IHC, IP, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

mouse, rat, human, canine

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse liver tissue, mouse lung tissue, mouse pancreas tissue

IP : mouse lung tissue,

IHC : human hepatocirrhosis tissue,

IF : HepG2 cells,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Shuang Xiao	34628270	Int Immunopharmacol	WB
Laura M Zidek	26113365	EMBO Rep	WB
Shun-Li Peng	35812404	Front Immunol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse liver tissue were subjected to SDS PAGE followed by western blot with 19495-1-AP (Aldehyde oxidase antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human hepatocirrhosis tissue slide using 19495-1-AP (Aldehyde oxidase antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human hepatocirrhosis tissue slide using 19495-1-AP (Aldehyde oxidase antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of HepG2 cells, using AOX1 antibody 19495-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red). Blue pseudocolor = DAPI (fluorescent DNA dye).



IP Result of anti-Aldehyde oxidase (IP:19495-1-AP, 4ug; Detection:19495-1-AP 1:300) with mouse lung tissue lysate 4000ug.