## For Research Use Only

## EIF2C2-Specific Polyclonal antibody

Catalog Number: 19691-1-AP

1 Publications



**Purification Method:** 

Antigen affinity purification

**Basic Information** 

Catalog Number:

19691-1-AP

150ul , Concentration: 133  $\mu$ g/ml by 27161 Bradford method using BSA as the

standard;

Source: Rabbit

Isotype:

GenBank Accession Number:

NM 012154 GeneID (NCBI):

**UNIPROT ID:** Q9UKV8

Full Name: eukaryotic translation initiation

factor 2C, 2 Calculated MW: 97 kDa

**Applications** 

**Tested Applications:** 

Cited Applications:

Species Specificity: human, mouse, rat Cited Species: human

**Background Information** 

EIF2C2, also named as AGO2, belongs to the argonaute family and Ago subfamily. It is required for RNA-mediated gene silencing (RNAi) by the RNA-induced silencing complex (RISC). The 'minimal RISC' appears to include EIF2C2/AGO2 bound to a short guide RNA such as a microRNA (miRNA) or short interfering RNA (siRNA). EIF2C2 may inhibit translation initiation by binding to the 7-methylguanosine cap, thereby preventing the recruitment of the translation initiation factor eIF4-E. It also inhibit translation initiation via interaction with EIF6, which itself binds to the 60S ribosomal subunit and prevents its association with the 40S ribosomal subunit. EIF2C2 can also  $upregulate \ the \ translation \ of \ specific \ mRNAs \ under \ certain \ growth \ conditions. \ This \ antibody \ is \ specific \ to \ EIF2C2.$ 

**Notable Publications** 

Author	Pubmed ID	Journal	Application
HuSheng Shan	37052807	Mol Biotechnol	RIP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA

Selected Validation Data