

For Research Use Only

# PAK2 Polyclonal antibody

Catalog Number: 19979-1-AP



## Basic Information

<b>Catalog Number:</b> 19979-1-AP	<b>GenBank Accession Number:</b> NM_002577	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 350 µg/ml by Nanodrop and 267 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 5062	<b>Recommended Dilutions:</b> WB 1:1000-1:5000 IF 1:50-1:500
<b>Source:</b> Rabbit	<b>Full Name:</b> p21 protein (Cdc42/Rac)-activated kinase 2	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 58 kDa	
	<b>Observed MW:</b> 54 kDa	

## Applications

<b>Tested Applications:</b> IF, WB, ELISA	<b>Positive Controls:</b> WB : HEK-293 cells, human skeletal muscle tissue, Jurkat cells, HeLa cells, PC-3 cells, Raji cells
<b>Species Specificity:</b> human	IF : HeLa cells,

## Background Information

PAK2, also named as PAK65, PAKgamma, p58, PAK-2p27, PAK-2p24 and C-t-PAK2, belongs to the protein kinase superfamily, STE Ser/Thr protein kinase family and STE20 subfamily. Full length PAK 2 stimulates cell survival and cell growth. The process is, at least in part, mediated by phosphorylation and inhibition of pro-apoptotic BAD. PAK2 has several isoforms with the MW of 54-62 kDa and 41 kDa. The 62 kDa PAK2 is cleaved into a 34 kDa C terminal fragment and a 28 kDa N terminal fragment with a time course that parallels apoptotic death in Jurkat cells. (PMID:10200518). Caspase-activated PAK-2p34 is involved in cell death response, probably involving the JNK signaling pathway. Cleaved PAK-2p34 seems to have a higher activity than the CDC42-activated form. The antibody has no cross reaction to PAK1 and PAK3.

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using PAK2 antibody (19979-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Untreated and Staurosporine treated Jurkat cells were subjected to SDS PAGE followed by western blot with 19979-1-AP (PAK2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 19979-1-AP (PAK2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.