For Research Use Only

MED12L Polyclonal antibody Catalog Number: 20047-1-AP



Basic Information	Catalog Number: 20047-1-AP	GenBank Accession Number: NM_053002	Purification Method: Antigen affinity purification
	Size: 150ul, Concentration: 300 ug/ml by Nanodrop and 333 ug/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG	GeneID (NCBI): 116931 UNIPROT ID:	Recommended Dilutions: WB 1:500-1:1000
		Q86YW9 Full Name: mediator complex subunit 12-like Calculated MW:	
		240 kDa Observed MW: 260-280 kDa	
Applications	Tested Applications: WB, ELISA Species Specificity: mouse	Positive Controls: WB : mouse liver tissue,	
Background Information	MED12L, also named as KIAA1635, TNRC11L, TRALP and TRALPUSH, belongs to the Mediator complex subunit 12 family. MED12L may be a component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. MED12L is a mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. MED12L is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. The antibody can recognize all the isoform of MED12L. The antibody has no cross reaction to MED12.		
Storage	Storage: Store at -20°C. Stable for one year aft Storage Buffer: PBS with 0.02% sodium azide and 50	% glycerol pH 7.3.	
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage	

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse liver tissue were subjected to SDS PAGE followed by western blot with 20047-1-AP (MED12L antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.