

For Research Use Only

SLC43A3 Polyclonal antibody

Catalog Number: 20131-1-AP



Basic Information

Catalog Number: 20131-1-AP	GenBank Accession Number: BC003163	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 550 ug/ml by Nanodrop;	GeneID (NCBI): 29015	Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500
Source: Rabbit	UNIPROT ID: Q8NBI5	
Isotype: IgG	Full Name: solute carrier family 43, member 3	
Immunogen Catalog Number: AG13999	Calculated MW: 491 aa, 55 kDa	
	Observed MW: 55 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls:
Species Specificity: human	WB : HL-60 cells, MCF-7 cells
	IHC : human ovary cancer tissue,

Note-IHC: suggested antigen retrieval with **TE buffer pH 9.0; (*)** Alternatively, antigen retrieval may be performed with **citrate buffer pH 6.0**

Background Information

Sodium-independent purine-selective nucleobase transporter(SLC43A3), also called as equilibrative nucleobase transporter 1(ENBT1), is a nucleobase transporter involved in purine salvage pathways in mammals, which mediates the equilibrative transport of extracellular purine nucleobases such as adenine, guanine and hypoxanthine.(PubMed:26455426, PubMed:32339528) It has been found that it may regulate the transport of fatty acids (FA) in fat cells, as a positive regulator of FA efflux and as a negative regulator of FA uptake.(PMID: 32217606)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 20131-1-AP (SLC43A3 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 20131-1-AP (SLC43A3 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).