For Research Use Only

ARF5-Specific Polyclonal antibody

Catalog Number:20227-1-AP

Featured Product 2 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 20227-1-AP	GenBank Accession Number: Purification Method: NM_001662 Antigen affinity purification GeneID (NCBI): Recommended Dilutions: by 381 WB 1:500-1:1000 UNIPROT ID: IHC 1:50-1:500 P84085 IF/ICC 1:50-1:500 Full Name: ADP-ribosylation factor 5 Calculated MW: 21 kDa					
	Size: 150ul , Concentration: 400 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG			WB 1:500-1:1000 IHC 1:50-1:500			
					Observed MW:		
				21 kDa			
				Applications	Tested Applications:	Positive Controls:	
		WB, IHC, IF/ICC, ELISA			WB : HepG2 cells, mouse colon tissue, mouse liver		
						ue, HeLa cells, mouse kidney tissue, mouse brain ue, COLO 320 cells, human testis tissue	
			stomach tissue, human skin cancer tissu				
human, mouse, rat			eLa cells,				
Cited Species: mouse			,				
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0							
Background Information	ADP-ribosylation factors (ARFs) are n 20kda protein size. ARFs bind and reg inactive GDP-bound conformations. A	gulate GTP/GDP cycle by NRF family proteins are envices and the second s Second second	alternating b ssential and o mmalian cell RF5 are class	ubiquitous in eukaryotes. Six highly .s. They function in vesicular traffic and II ARFs. (PMID: 7759471, PMID:			
	ADP-ribosylation factors (ARFs) are n 20kda protein size. ARFs bind and reg inactive GDP-bound conformations. A conserved members of the family ha actin remodelling and other bioproce 16042562). Their function are not full	gulate GTP/GDP cycle by NRF family proteins are envices and the second s Second second	alternating b ssential and i mmalian cell RF5 are class / is specific to	etween the active GTP-bound and ubiquitous in eukaryotes. Six highly s. They function in vesicular traffic and II ARFs. (PMID: 7759471, PMID:			
Background Information	ADP-ribosylation factors (ARFs) are n 20kda protein size. ARFs bind and reg inactive GDP-bound conformations. A conserved members of the family ha actin remodelling and other bioproce 16042562). Their function are not full Author Put	gulate GTP/GDP cycle by NF family proteins are e- ve been identified in ma isses in cells. ARF4 and A y described.The antibody	alternating b ssential and (mmalian cell RF 5 are class / is specific to	etween the active GTP-bound and ubiquitous in eukaryotes. Six highly .s. They function in vesicular traffic and II ARFs. (PMID: 7759471, PMID: 5 ARF5.			
	ADP-ribosylation factors (ARFs) are n 20kda protein size. ARFs bind and reg inactive GDP-bound conformations. A conserved members of the family ha actin remodelling and other bioproce 16042562). Their function are not full Author Put Rakhee Bajaj 36:	gulate GTP/GDP cycle by IRF family proteins are en- ve been identified in ma- sses in cells. ARF4 and A y described. The antibody pmed ID Journal	alternating b ssential and (mmalian cell RF 5 are class / is specific to	etween the active GTP-bound and ubiquitous in eukaryotes. Six highly s. They function in vesicular traffic and II ARFs. (PMID: 7759471, PMID: o ARF5. Application			

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 20227-1-AP (ARF5-Specific antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human stomach tissue slide using 20227-1-AP (ARF5-Specific antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded skin cancer slide using 20227-1-AP (ARF5-Specific antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 20227-1-AP (ARF5-Specific antibody) at dilution of 1:50 and Alexa Fluor 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).